



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01H 1/00, 5/00, 5/10, C12N 15/29, 5/04, 5/10	A1	(11) International Publication Number: WO 99/41974 (43) International Publication Date: 26 August 1999 (26.08.99)
(21) International Application Number: PCT/US99/03429 (22) International Filing Date: 17 February 1999 (17.02.99) (30) Priority Data: 09/026,039 19 February 1998 (19.02.98) US (71) Applicant (for all designated States except US): THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 12th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): JOFUKU, K., Diane [US/US]; 475 Sims Road, Santa Cruz, CA 95060 (US). OKAMURO, Jack, K. [US/US]; 475 Sims Road, Santa Cruz, CA 95060 (US). (74) Agents: BASTIAN, Kevin, L. et al.; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111-3834 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHODS FOR IMPROVING SEEDS (57) Abstract The invention provides methods of modulating seed mass and other traits in plants. The methods involve producing transgenic plants comprising a recombinant expression cassette containing an ADC nucleic acid linked to a plant promoter.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS FOR IMPROVING SEEDS

REFERENCE TO RELATED APPLICATIONS

This is a continuation in part of copending application USSN 08/912,272, which is a continuation in part of USSN 08/879,827, which is a continuation in part of USSN 08/700,152, filed August 20, 1996, all of which are incorporated herein by reference.

FIELD OF THE INVENTION

The present invention is directed to plant genetic engineering. In particular, it relates to new methods for modulating mass and other properties of plant seeds.

BACKGROUND OF THE INVENTION

The pattern of flower development is controlled by the floral meristem, a complex tissue whose cells give rise to the different organ systems of the flower. Genetic and molecular studies have defined an evolutionarily conserved network of genes that control floral meristem identity and floral organ development in *Arabidopsis*, snapdragon, and other plant species (*see, e.g.,* Coen and Carpenter, *Plant Cell* 5:1175-1181 (1993) and Okamuro *et al.*, *Plant Cell* 5:1183-1193 (1993)). In *Arabidopsis*, a floral homeotic gene *APETALA2* (*AP2*) controls three critical aspects of flower ontogeny - the establishment of the floral meristem (Irish and Sussex, *Plant Cell* 2:741-753 (1990); Huala and Sussex, *Plant Cell* 4:901-913 (1992); Bowman *et al.*, *Development* 119:721-743 (1993); Schultz and Haughn, *Development* 119:745-765 (1993); Shannon and Meeks-Wagner, *Plant Cell* 5:639-655 (1993)), the specification of floral organ identity (Komaki *et al.*, *Development* 104:195-203 (1988)); Bowman *et al.*, *Plant Cell* 1:37-52 (1989); Kunst *et al.*, *Plant Cell* 1:1195-1208 (1989)), and the temporal and spatial regulation of floral homeotic gene expression (Bowman *et al.*, *Plant Cell* 3:749-758 (1991); Drews *et al.*, *Cell* 65:91-1002 (1991)).

One early function of *AP2* during flower development is to promote the establishment of the floral meristem. *AP2* performs this function in cooperation with at

least three other floral meristem genes, *APETALA1* (*API*), *LEAFY* (*LFY*), and *CAULIFLOWER* (*CAL*) (Irish and Sussex (1990); Bowman, *Flowering Newsletter* 14:7-19 (1992); Huala and Sussex (1992); Bowman *et al.*, (1993); Schultz and Haughn, (1993); Shannon and Meeks-Wagner, (1993)). A second function of *AP2* is to regulate floral organ development. In *Arabidopsis*, the floral meristem produces four concentric rings or whorls of floral organs - sepals, petals, stamens, and carpels. In weak, partial loss-of-function *ap2* mutants, sepals are homeotically transformed into leaves, and petals are transformed into pollen-producing stamenoid organs (Bowman *et al.*, *Development* 112:1-20 (1991)). By contrast, in strong *ap2* mutants, sepals are transformed into ovule-bearing carpels, petal development is suppressed, the number of stamens is reduced, and carpel fusion is often defective (Bowman *et al.*, (1991)). Finally, the effects of *ap2* on floral organ development are in part a result of a third function of *AP2*, which is to directly or indirectly regulate the expression of several flower-specific homeotic regulatory genes (Bowman *et al.*, *Plant Cell* 3:749-758 (1991); Drews *et al.*, *Cell* 65:91-1002 (1991); Jack *et al.* *Cell* 68:683-697 (1992); Mandel *et al.* *Cell* 71: 133-143 (1992)).

Clearly, *Ap2* plays a critical role in the regulation of *Arabidopsis* flower development. Yet, little is known about how it carries out its functions at the cellular and molecular levels. A spatial and combinatorial model has been proposed to explain the role of *AP2* and other floral homeotic genes in the specification of floral organ identity(see, e.g., Coen and Carpenter, *supra*). One central premise of this model is that *AP2* and a second floral homeotic gene *AGAMOUS* (*AG*) are mutually antagonistic genes. That is, *AP2* negatively regulates *AG* gene expression in sepals and petals, and conversely, *AG* negatively regulates *AP2* gene expression in stamens and carpels. In situ hybridization analysis of *AG* gene expression in wild-type and *ap2* mutant flowers has demonstrated that *AP2* is indeed a negative regulator of *AG* expression. However, it is not yet known how *AP2* controls *AG*. Nor is it known how *AG* influences *AP2* gene activity.

The *AP2* gene in *Arabidopsis* has been isolated by T-DNA insertional mutagenesis as described in Jofuku *et al.* *The Plant Cell* 6:1211-1225 (1994). *AP2* encodes a putative nuclear factor that bears no significant similarity to any known fungal, or animal regulatory protein. Evidence provided there indicates that *AP2* gene activity

and function are not restricted to developing flowers, suggesting that it may play a broader role in the regulation of *Arabidopsis* development than originally proposed.

In spite of the recent progress in defining the genetic control of plant development, little progress has been reported in the identification and analysis of genes effecting agronomically important traits such as seed size, protein content, oil content and the like. Characterization of such genes would allow for the genetic engineering of plants with a variety of desirable traits. The present invention addresses these and other needs.

SUMMARY OF THE INVENTION

The present invention provides methods of modulating seed mass and other traits in plants. The methods involve providing a plant comprising a recombinant expression cassette containing an *ADC* nucleic acid linked to a plant promoter. The plant is either selfed or crossed with a second plant to produce a plurality of seeds. Seeds with the desired trait (*e.g.*, altered mass) are then selected.

In some embodiments, transcription of the *ADC* nucleic acid inhibits expression of an endogenous *ADC* gene or activity the encoded protein. In these embodiments, the step of selecting includes the step of selecting seed with increased mass or another trait. The seed may have, for instance, increased protein content, carbohydrate content, or oil content. In the case of increased oil content, the types of fatty acids may or may not be altered as compared to the parental lines. In these embodiments, the *ADC* nucleic acid may be linked to the plant promoter in the sense or the antisense orientation. Alternatively, expression of the *ADC* nucleic acid may enhance expression of an endogenous *ADC* gene or *ADC* activity and the step of selecting includes the step of selecting seed with decreased mass. This embodiment is particularly useful for producing seedless varieties of crop plants.

If the first plant is crossed with a second plant the two plants may be the same or different species. The plants may be any higher plants, for example; members of the families Brassicaceae or Solanaceae. In making seed of the invention, either the female or the male parent plant can comprise the expression cassette containing the *ADC* nucleic acid. In preferred embodiments, both parents contain the expression cassette.

In the expression cassettes, the plant promoter may be a constitutive promoter, for example, the CaMV 35S promoter. Alternatively, the promoter may be a

tissue-specific promoter. Examples of tissue specific expression useful in the invention include fruit-specific, seed-specific (e.g., ovule-specific, embryo-specific, endosperm-specific, integument-specific, or seed coat-specific) expression.

The invention also provides seed produced by the methods described above. The seed of the invention comprise a recombinant expression cassette containing an *ADC* nucleic acid. If the expression cassette is used to inhibit expression of endogenous *ADC* expression, the seed will have a mass at least about 20% greater than the average mass of seeds of the same plant variety which lack the recombinant expression cassette. If the expression cassette is used to enhance expression of *ADC*, the seed will have a mass at least about 20% less than the average mass of seeds of the same plant variety which lack the recombinant expression cassette. Other traits such as protein content, carbohydrate content, and oil content can be altered in the same manner.

Definitions

The phrase "nucleic acid sequence" refers to a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. It includes chromosomal DNA, self-replicating plasmids, infectious polymers of DNA or RNA and DNA or RNA that performs a primarily structural role..

The term "promoter" refers to a region or sequence determinants located upstream or downstream from the start of transcription and which are involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells.

The term "plant" includes whole plants, plant organs (e.g., leaves, stems, flowers, roots, etc.), seeds and plant cells and progeny of same. The class of plants which can be used in the method of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants), as well as gymnosperms. It includes plants of a variety of ploidy levels, including polyploid, diploid, haploid and hemizygous.

A polynucleotide sequence is "heterologous to" an organism or a second polynucleotide sequence if it originates from a foreign species, or, if from the same species, is modified from its original form. For example, a promoter operably linked to a heterologous coding sequence refers to a coding sequence from a species different from

that from which the promoter was derived, or, if from the same species, a coding sequence which is different from any naturally occurring allelic variants. As defined here, a modified *ADC* coding sequence which is heterologous to an operably linked *ADC* promoter does not include the T-DNA insertional mutants (*e.g.*, *ap2-10*) as described in Jofuku *et al. The Plant Cell* 6:1211-1225 (1994).

A polynucleotide "exogenous to" an individual plant is a polynucleotide which is introduced into the plant by any means other than by a sexual cross. Examples of means by which this can be accomplished are described below, and include *Agrobacterium*-mediated transformation, biolistic methods, electroporation, and the like. Such a plant containing the exogenous nucleic acid is referred to here as an R_1 generation transgenic plant. Transgenic plants which arise from sexual cross or by selfing are descendants of such a plant.

An "ADC (AP2 domain containing) nucleic acid" or "ADC polynucleotide sequence" of the invention is a subsequence or full length polynucleotide sequence of a gene which, encodes an polypeptide containing an AP2 domain and when present in a transgenic plant, can be used to modulate seed properties in seed produced by the plant. An exemplary nucleic acid of the invention is the *Arabidopsis AP2* sequence as disclosed in Jofuku *et al. The Plant Cell* 6:1211-1225 (1994). The GenBank accession number for this sequence is U12546. As explained in detail below a family of *RAP2* (related to *AP2*) genes have been identified in *Arabidopsis*. The class of nucleic acids claimed here falls into at least two subclasses (AP2-like and EREBP-like genes), which are distinguished by, for instance, the number of AP2 domains contained within each polypeptide and by sequences within certain conserved regions. The differences between these two subclasses are described in more detail below. *ADC* polynucleotides are defined by their ability to hybridize under defined conditions to the exemplified nucleic acids or PCR products derived from them. An *ADC* polynucleotide (*e.g.*, *AP2* or *RAP2*) is typically at least about 30-40 nucleotides to about 3000, usually less than about 5000 nucleotides in length. Usually the nucleic acids are from about 100 to about 2000 nucleotides, often from about 500 to about 1700 nucleotides in length.

ADC nucleic acids, as explained in more detail below, are a new class of plant regulatory genes that encode *ADC* polypeptides, which are distinguished by the presence of one or more of a 56-68 amino acid repeated motif, referred to here as the "AP2 domain". The amino acid sequence of an exemplary AP2 polypeptide is shown in

Jofuku *et al.*, *supra*. One of skill will recognize that in light of the present disclosure various modifications (*e.g.*, substitutions, additions, and deletions) can be made to the sequences shown there without substantially affecting its function. These variations are specifically covered by the terms ADC polypeptide or ADC polynucleotide.

5 In the case of both expression of transgenes and inhibition of endogenous genes (*e.g.*, by antisense, or sense suppression) one of skill will recognize that the inserted polynucleotide sequence need not be identical, but may be only "substantially identical" to a sequence of the gene from which it was derived. As explained below, these substantially identical variants are specifically covered by the term ADC nucleic acid.

10 In the case where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, one of skill will recognize that because of codon degeneracy a number of polynucleotide sequences will encode the same polypeptide. These variants are specifically covered by the terms "ADC nucleic acid", "AP2 nucleic acid" and "RAP2 nucleic acid". In addition, the term specifically includes those full length sequences substantially identical (determined as described below) with an ADC polynucleotide sequence and that encode proteins that retain the function of the ADC polypeptide (*e.g.*, resulting from conservative substitutions of amino acids in the AP2 polypeptide). In addition, variants can be those that encode dominant negative mutants as described below.

15 Two nucleic acid sequences or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the complementary sequence is identical to all or a portion of a reference polynucleotide sequence.

20 Sequence comparisons between two (or more) polynucleotides or polypeptides are typically performed by comparing sequences of the two sequences over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window", as used herein, refers to a segment of at least about 20 contiguous positions, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

"Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 60% sequence identity, preferably at least 80%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using the programs described above (preferably BLAST) using standard parameters. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 35%, preferably at least 60%, more preferably at least 90%, and most preferably at least 95%. Polypeptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side

chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other, or a third nucleic acid, under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least about 60°C.

In the present invention, genomic DNA or cDNA comprising *ADC* nucleic acids of the invention can be identified in standard Southern blots under stringent conditions using the nucleic acid sequences disclosed here. For the purposes of this disclosure, stringent conditions for such hybridizations are those which include at least one wash in 0.2X SSC at a temperature of at least about 50°C, usually about 55°C to about 60°C, for 20 minutes, or equivalent conditions. Other means by which nucleic acids of the invention can be identified are described in more detail below.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A shows amino acid sequence alignment between AP2 direct repeats AP2-R1 (aa 129-195) and AP2-R2 (aa 221-288). Solid and dashed lines between the two sequences indicate residue identity and similarity, respectively. Arrows indicate the positions of the ap2-1, ap2-5, and ap2-10 mutations described in Jofuku *et al.* (1994). The bracket above the AP2-R1 and AP2-R2 sequences indicates the residues capable of forming amphipathic α -helices shown in Figure 1B.

Figure 1B is a schematic diagram of the putative AP2-R1 (R1) and AP2-R2 (R2) amphipathic α -helices. The NH₂ terminal ends of the R1 and R2 helices begin

at residues Phe-160 and Phe-253 and rotate clockwise by 100° per residue through Phe-177 and Cys-270, respectively. Arrows directed toward or away from the center of the helical wheel diagrams indicate the negative or positive degree of hydrophobicity as defined by Jones *et al.* *J. Lipid Res.* 33: 87-296 (1992).

5 Figure 2 shows an antisense construct of the invention. pPW14.4 (which is identical to pPW15) represents the 13.41 kb AP2 antisense gene construct used in plant transformation described here. pPW14.4 is comprised of the AP2 gene coding region in a transcriptional fusion with the cauliflower mosaic virus 35S (P35S) constitutive promoter in an antisense orientation. The Ti plasmid vector used is a
10 modified version of the pGSJ780A vector (Plant Genetic Systems, Gent, Belgium) in which a unique EcoR1 restriction site was introduced into the BamH1 site using a Cla1-EcoR1-BamH1 adaptor. The modified pGSJ780A vector DNA was linearized with EcoR1 and the AP2 coding region inserted as a 1.68 kb EcoR1 DNA fragment from AP2 cDNA plasmid cAP2#1 (Jofuku *et al.*, 1994) in an antisense orientation with respect to
15 the 35S promoter. KmR represents the plant selectable marker gene NPTII which confers resistance to the antibiotic kanamycin to transformed plant cells carrying an integrated 35S-AP2 antisense gene. Boxes 1 and 5 represent the T-DNA left and right border sequences, respectively, that are required for transfer of T-DNA containing the 35S-AP2 antisense gene construct into the plant genome. Regions 2 and 3 contain T-
20 DNA sequences. Box 3 designates the 3' octopine synthase gene sequences that function in transcriptional termination. Region 6 designates bacterial DNA sequences that function as a bacterial origin of replication in both *E. coli* and *Agrobacterium tumefaciens*, thus allowing pPW14.4 plasmid replication and retention in both bacteria. Box 7 represents the bacterial selectable marker gene that confers resistance to the
25 antibiotics streptomycin and spectinomycin and allows for selection of *Agrobacterium* strains that carry the pPW14.4 recombinant plasmid.

 Figure 3 shows a sense construct of the invention. pPW12.4 (which is identical to pPW9) represents the 13.41 kb AP2 sense gene construct used in plant transformation described here. pPW12.4 is comprised of the AP2 gene coding region in
30 a transcriptional fusion with the cauliflower mosaic virus 35S (P35S) constitutive promoter in a sense orientation. The Ti plasmid vector used is a modified version of the pGSJ780A vector (Plant Genetic Systems, Gent, Belgium) in which a unique EcoR1 restriction site was introduced into the BamH1 site using a Cla1-EcoR1-BamH1 adaptor.

The modified pGSJ780A vector DNA was linearized with EcoR1 and the AP2 coding region inserted as a 1.68 kb EcoR1 DNA fragment from AP2 cDNA plasmid cAP2#1 (Jofuku et al., 1994) in a sense orientation with respect to the 35S promoter. KmR represents the plant selectable marker gene NPTII which confers resistance to the antibiotic kanamycin to transformed plant cells carrying an integrated 35S-AP2 antisense gene. Boxes 1 and 5 represent the T-DNA left and right border sequences, respectively, that are required for transfer of T-DNA containing the 35S-AP2 sense gene construct into the plant genome. Regions 2 and 3 contain T-DNA sequences. Box 3 designates the 3' octopine synthase gene sequences that function in transcriptional termination. Region 6 designates bacterial DNA sequences that function as a bacterial origin of replication in both *E. coli* and *Agrobacterium tumefaciens*, thus allowing pPW12.4 plasmid replication and retention in both bacteria. Box 7 represents the bacterial selectable marker gene that confers resistance to the antibiotics streptomycin and spectinomycin and allows for selection of *Agrobacterium* strains that carry the pPW12.4 recombinant plasmid.

Figures 4A and 4B show AP2 domain sequence and structure. The number of amino acid residues within each AP2 domain is shown to the right. Sequence gaps were introduced to maximize sequence alignments. The position of amino acid residues and sequence gaps within the AP2 domain alignments are numbered 1-77 for reference. The location of the conserved YRG and RAYD elements are indicated by brackets. Shaded boxes highlight regions of sequence similarity. Positively charged amino acids within the YRG element are indicated by + signs above the residues. The location of the 18-amino acid core region that is predicted to form an amphipathic α -helix in AP2 is indicated by a bracket. Residues within the RAYD element of each AP2 domain that are predicted to form an amphipathic α -helix are underlined. Figure 4A shows members of the AP2-like subclass. Amino acid sequence alignment between the AP2 domain repeats R1 and R2 contained within AP2, ANT and RAP2.7 is shown. Brackets above the sequences designate the conserved YRG and RAYD blocks described above. The filled circle and asterisk indicate the positions of the *ap2-1*, and *ap2-5* mutations, respectively. Amino acid residues that constitute a consensus AP2 domain motif for AP2, ANT, and RAP2.7 is shown below the alignment with invariant residues shown capitalized. Figure 4B shows members of the EREBP-like subclass. Amino acid sequence alignment between the AP2 domains contained within the tobacco EREBPs and the *Arabidopsis* EREBP-like RAP2 proteins is shown. GenBank accession numbers for

EREBP-1, EREBP-2, EREBP-3, and EREBP-4 are D38123, D38126, D38124, and D38125, respectively.

Figure 4C provides schematic diagrams of the putative RAP2.7-R1, AP2-R1, and ANT-R1 amphipathic α -helices. Amino acid residues within the RAP2.7-R1, AP2-R1, and ANT-R1 motifs shown underlined in A that are predicted to form amphipathic α -helices are schematically displayed with residues rotating clockwise by 100° per residue to form helical structures. Arrows directed toward or away from the center of the helical wheel diagrams indicate the negative or positive degree of hydrophobicity as defined by Jones *et al. J. Lipid Res.* 33:287-296 (1992). Positively and negatively charged amino acid residues are designated by + and - signs, respectively.

Figure 4D shows schematic diagrams of the putative RAP2.2, RAP2.5, RAP2-12, and EREBP-3 amphipathic α -helices. Amino acid residues within the RAP2.2, RAP2.5, RAP2-12, and EREBP-3 motifs shown underlined in Figure 4B that are predicted to form amphipathic α -helices are schematically displayed as described in Figure 4C.

Figure 4E shows sequence alignment between the 25-26 amino acid linker regions in AP2, ANT, and RAP2.7. R1 and R2 designate the positions of the R1 and R2 repeats within AP2, ANT, and RAP2.7 relative to the linker region sequences. Boxes designate invariant residues within the conserved linker regions. Amino acid residues that constitute a consensus linker region motif for AP2, ANT, and RAP2.7 are shown below the alignment with invariant residues shown capitalized. The arrowhead indicates the position of the *ant-3* mutation described by Klucher *et al. Plant Cell* 8:137-153 (1996).

Figure 5 is a schematic diagram of pAP2, which can be used to construct expression vectors of the invention.

Figure 6 is a schematic diagram of pBEL1, which can be used to construct expression vectors of the invention.

Figure 7 is schematic diagram of gene expression.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

This invention relates to plant *ADC* genes, such as the *AP2* and *RAP2* genes of *Arabidopsis*. The invention provides molecular strategies for controlling seed size and total seed protein using *ADC* overexpression and antisense gene constructs. In particular, transgenic plants containing antisense constructs have dramatically increased seed mass, seed protein, or seed oil. Alternatively, overexpression of *ADC* using a constructs of the invention leads to reduced seed size and total seed protein. Together, data presented here demonstrate that a number of agronomically important traits including seed mass, total seed protein, and oil content, can be controlled in species of agricultural importance.

Isolation of *ADC* nucleic acids

Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's specifications. These techniques and various other techniques are generally performed according to Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989).

The isolation of *ADC* nucleic acids may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed here can be used to identify the desired gene in a cDNA or genomic DNA library. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a cDNA library, mRNA is isolated from the desired organ, such as flowers, and a cDNA library which contains the *ADC* gene transcript is prepared from the mRNA. Alternatively, cDNA may be prepared from mRNA extracted from other tissues in which *ADC* genes or homologs are expressed.

The cDNA or genomic library can then be screened using a probe based upon the sequence of a cloned *ADC* gene disclosed here. Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the

same or different plant species. Alternatively, antibodies raised against an ADC polypeptide can be used to screen an mRNA expression library.

Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of the *ADC* genes directly from genomic DNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes.

Appropriate primers and probes for identifying *ADC* sequences from plant tissues are generated from comparisons of the sequences provided in Jofuku *et al.*, *supra*. For a general overview of PCR see *PCR Protocols: A Guide to Methods and Applications*. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), *Academic Press*, San Diego (1990).

As noted above, the nucleic acids of the invention are characterized by the presence of sequence encoding a AP2 domain. Thus, these nucleic acids can be identified by their ability to specifically hybridize to sequences encoding AP2 domain disclosed here. Primers which specifically amplify AP2 domains of the exemplified genes are particularly useful for identification of particular *ADC* polynucleotides. Primers suitable for this purpose based on the sequence of RAP2 genes disclosed here are as follows:

Name	GenBank Number	Primers
AP2	U12546	JOAP2U 5'-GTTGCCGCTGCCGTAGTG-3' JOAP2L 5'-GGTTCATCCTGAGCCGCATATC-3'
RAP2.1	AF003094	JORAP2.1U 5'-CTCAAGAAGAAGTGCCTAACCACG-3' JORAP2.1L 5'-GCAGAAGCTAGAAGAGCGTCGA-3'
RAP2.2	AF003095	JORAP2.2U 5'-GGAAAATGGGCTGCGGAG-3' JORAP2.2L 5'-GTTACCTCCAGCATCGAACGAG-3'
RAP2.4	AF003097	JORAP2.4U 5'-GCTGGATCTTGTTTCGCTTACG-3' JORAP2.4L 5'-GCTTCAAGCTTAGCGTCGACTG-3'
RAP2.5	AF003098	JORAP2.5U 5'-AGATGGGCTTGAAACCCGAC-3' JORAP2.5L 5'-CTGGCTAGGGCTACGCGC-3'
RAP2.6	AF003099	JORAP2.6U 5'-TTCTTTGCCTCCTCAACCATTG-3' JORAP2.6L 5'-TCTGAGTTCCAACATTTTCGGG-3'
RAP2.7	AF003100	JORAP2.7U 5'-GAAATTGGTAACTCCGGTTCCG-3' JORAP2.7L 5'-CCATTTTGCTTTGGCGCATTAC-3'
RAP2.8	AF003101	JORAP2.8U 5'-GGCGTTACGCCTCTACCGG-3' JORAP2.8L 5'-CGCCGTCTTCCAGAACGTTC-3'
RAP2.9	AF003102	JORAP2.9U 5'-ATCACGGATCTGGCTTGGTTC-3' JORAP2.9L 5'-GCCTTCTTCCGTATCAACGTTCG-3'
RAP2.10	AF003103	JORAP2.10U 5'-GTCAACTCCGGCGGTTACG-3' JORAP2.10L 5'-TCTCCTTATATACGCCGCCGA-3'
RAP2.11	AF003104	JORAP2.11U 5'-GAGAAGAGCAAAGGCAACAAGAC-3' JORAP2.11L 5'-AGTTGTTAGGAAAATGGTTTGCG-3'
RAP2.12	AF003105	JORAP2.12U 5'-AAACCATTCGTTTTCACTTCGACTC-3' JORAP2.12L 5'-TCACAGAGCGTTTCTGAGAATTAGC-3'

The PCR primers are used under standard PCR conditions (described for instance in Innis *et al.*) using the nucleic acids as identified in the above GenBank accessions as a template. The PCR products generated by any of the reactions can then be used to identify nucleic acids of the invention (*e.g.*, from a cDNA library) by their ability to hybridize to these products. Particularly preferred hybridization conditions use a Hybridization Buffer consisting of: 0.25M Phosphate Buffer (pH 7.2), 1 mM EDTA, 1% Bovine Serum Albumin, 7% SDS. Hybridizations then followed by a first wash with 2.0XSSC + 0.1% SDS or 0.39M Na⁺ and subsequent washes with 0.2XSSC + 0.1% SDS or 0.042M Na⁺. Hybridization temperature will be from about 45°C to about 78°C, usually from about 50°C to about 70°C. Followed by washes at 18°C.

Particularly preferred hybridization conditions are as follows:

Hybridization Temp.	Hybrid. Time	Wash Buffer A	Wash Buffer B
78 degrees C	48 hrs	18 degrees C	18 degrees C
70 degrees C	48 hrs	18 degrees C	18 degrees C
65 degrees C	48 hrs	18 degrees C	18 degrees C
60 degrees C	72 hrs	18 degrees C	18 degrees C
55 degrees C	96 hrs	18 degrees C	18 degrees C
45 degrees C	200 hrs	18 degrees C	No wash

If desired, primers that amplify regions more specific to particular ADC genes can be used. The PCR products produced by these primers can be used in the hybridization conditions described above to isolate nucleic acids of the invention.

Name	GenBank Number	Primers
AP2	U12546	AP2U 5'-ATGTGGGATCTAAACGACGCAC-3' AP2L 5'-GATCTTGGTCCACGCCGAC-3'
RAP2.1	AF003094	RAP2.1U 5'-AAG AGG ACC ATC TCT CAG-3' RAP2.1L 5'-AAC ACT CGC TAG CTT CTC-3'
RAP2.2	AF003095	RAP2.2U 5'-TGG TTC AGC AGC CAA CAC-3' RAP2.2L 5'-CAA TGC ATA GAG CTT GAG G-3'

RAP2.4	AF003097	RAP2.4U 5'-ACG GAT TTC ACA TCG GAG-3' RAP2.4L 5'-CTA AGC TAG AAT CGA ATC C-3'
RAP2.5	AF003098	RAP2.5U 5'-TACCGGTTTCGCGCGTAG-3' RAP2.5L 5'-CACCTTCGAAATCAACGACCG-3'
RAP2.6	AF003099	RAP2.6U 5'-TTCCCCGAAAATGTTGGAAGTC-3' RAP2.6L 5'-TGGGAGAGAAAAAATTGGTAGATCG-3'
RAP2.7	AF003100	RAP2.7U 5'- CGA TGG AGA CGA AGA CTC-3' RAP2.7L 5'- GTC GGA ACC GGA GTT ACC-3'
RAP2.8	AF003101	RAP2.8U 5'-TCA CTC AAA GGC CGA GAT C-3' RAP2.8L 5'-TAA CAA CAT CAC CGG CTC G-3'
RAP2.9	AF003102	RAP2.9U 5'-GTG AAG GCT TAG GAG GAG-3' RAP2.9L 5'-TGC CTC ATA TGA GTC AGA G-3'
RAP2.10	AF003103	RAP2.10U 5'-TCCCGGAGCTTTTAGCCG-3' RAP2.10L 5'-CAACCCGTTCCAACGATCC-3'
RAP2.11	AF003104	RAP2.11U 5'-TTCTTCACCAGAAGCAGAGCATG-3' RAP2.11L 5'-CTCCATTCATTGCATATAGGGACG-3'
RAP2.12	AF003105	RAP2.12U 5'-GCTTTGGTTCAGAACTCGAACATC-3' RAP2.12L 5'-AGGTTGATAAACGAACGATGCG-3'

Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, e.g., Carruthers *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 47:411-418 (1982), and Adams *et al.*, *J. Am. Chem. Soc.* 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Alternatively, primers that specifically hybridize to highly conserved regions in AP2 domains can be used to amplify sequences from widely divergent plant species such as *Arabidopsis*, canola, soybean, tobacco, and snapdragon. Examples of such primers are as follows:

Primer RISZU 1: 5'-GGAYTGTGGGAAACAAGTTTA-3'

Primer RISZU 2: 5'-TGCAAAGTRACACCTCTATACTT-3'

Y = pyrimidine (T or C)

R = purine (A or G).

5 Standard nucleic acid hybridization techniques using the conditions disclosed above can then be used to identify full length cDNA or genomic clones.

In addition, the following DNA primers, RISZU 3 and RISZU 4, can be used in an inverse PCR reaction to specifically amplify flanking AP2 gene sequences from widely divergent plant species. These primers are as follows:

10 Primer RISZU 3: 5'-GCATGWGCAGTGTCAAATCCA-3'

Primer RISZU 4: 5'-GAGGAAGTTCVAAGTATAGA-3'

W = A or T

V = G, A, or C

15 These primers have been used in standard PCR conditions to amplify ADC gene sequences from canola (SEQ ID NO:1) and soybean (SEQ ID NO:2).

Control of ADC activity or gene expression

20 One of skill will recognize that a number of methods can be used to modulate ADC activity or gene expression. ADC activity can be modulated in the plant cell at the gene, transcriptional, posttranscriptional, translational, or posttranslational, levels as schematically shown in Figure 7. Techniques for modulating ADC activity at each of these levels are generally well known to one of skill and are discussed briefly below.

25 Methods for introducing genetic mutations into plant genes are well known. For instance, seeds or other plant material can be treated with a mutagenic chemical substance, according to standard techniques. Such chemical substances include, but are not limited to, the following: diethyl sulfate, ethylene imine, ethyl methanesulfonate and N-nitroso-N-ethylurea. Alternatively, ionizing radiation from
30 sources such as, for example, X-rays or gamma rays can be used. Desired mutants are selected by assaying for increased seed mass, oil content and other properties.

Alternatively, homologous recombination can be used to induce targeted gene disruptions by specifically deleting or altering the *ADC* gene *in vivo* (see, generally,

Grewal and Klar, *Genetics* 146: 1221-1238 (1997) and Xu *et al.*, *Genes Dev.* 10: 2411-2422 (1996)). Homologous recombination has been demonstrated in plants (Puchta *et al.*, *Experientia* 50: 277-284 (1994), Swoboda *et al.*, *EMBO J.* 13: 484-489 (1994); and Offringa *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 7346-7350 (1993)).

5 In applying homologous recombination technology to the genes of the invention, mutations in selected portions of an *ADC* gene sequences (including 5' upstream, 3' downstream, and intragenic regions) such as those disclosed here are made *in vitro* and then introduced into the desired plant using standard techniques. Since the efficiency of homologous recombination is known to be dependent on the vectors used,
10 use of dicistronic gene targeting vectors as described by Mountford *et al.* *Proc. Natl. Acad. Sci. USA* 91: 4303-4307 (1994); and Vaulont *et al.* *Transgenic Res.* 4: 247-255 (1995) are conveniently used to increase the efficiency of selecting for altered *ADC* gene expression in transgenic plants. The mutated gene will interact with the target wild-type gene in such a way that homologous recombination and targeted replacement of the wild-
15 type gene will occur in transgenic plant cells, resulting in suppression of *ADC* activity.

Alternatively, oligonucleotides composed of a contiguous stretch of RNA and DNA residues in a duplex conformation with double hairpin caps on the ends can be used. The RNA/DNA sequence is designed to align with the sequence of the target *ADC* gene and to contain the desired nucleotide change. Introduction of the chimeric
20 oligonucleotide on an extrachromosomal T-DNA plasmid results in efficient and specific *ADC* gene conversion directed by chimeric molecules in a small number of transformed plant cells. This method is described in Cole-Strauss *et al.* *Science* 273:1386-1389 (1996) and Yoon *et al.* *Proc. Natl. Acad. Sci. USA* 93: 2071-2076 (1996).

Gene expression can be inactivated using recombinant DNA techniques by
25 transforming plant cells with constructs comprising transposons or T-DNA sequences. *ADC* mutants prepared by these methods are identified according to standard techniques. For instance, mutants can be detected by PCR or by detecting the presence or absence of *ADC* mRNA, *e.g.*, by Northern blots. Mutants can also be selected by assaying for increased seed mass, oil content and other properties.

30 The isolated nucleic acid sequences prepared as described herein, can also be used in a number of techniques to control endogenous *ADC* gene expression at various levels. Subsequences from the sequences disclosed here can be used to control, transcription, RNA accumulation, translation, and the like.

A number of methods can be used to inhibit gene expression in plants. For instance, antisense technology can be conveniently used. To accomplish this, a nucleic acid segment from the desired gene is cloned and operably linked to a promoter such that the antisense strand of RNA will be transcribed. The construct is then
5 transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been suggested that antisense suppression can act at all levels of gene regulation including suppression of RNA translation (*see*, Bourque *Plant Sci. (Limerick)* 105: 125-149 (1995); Pantopoulos *In Progress in Nucleic Acid Research and Molecular Biology*, Vol. 48. Cohn, W. E. and K. Moldave (Ed.). Academic Press, Inc.: San Diego,
10 California, USA; London, England, UK. p. 181-238; Heiser *et al. Plant Sci. (Shannon)* 127: 61-69 (1997)) and by preventing the accumulation of mRNA which encodes the protein of interest, (*see*, Baulcombe *Plant Mol. Bio.* 32:79-88 (1996); Prins and Goldbach *Arch. Virol.* 141: 2259-2276 (1996); Metzlauff *et al. Cell* 88: 845-854 (1997), Sheehy *et al., Proc. Nat. Acad. Sci. USA*, 85:8805-8809 (1988), and Hiatt *et al., U.S.*
15 Patent No. 4,801,340).

The nucleic acid segment to be introduced generally will be substantially identical to at least a portion of the endogenous *ADC* gene or genes to be repressed. The sequence, however, need not be perfectly identical to inhibit expression. The vectors of the present invention can be designed such that the inhibitory effect applies to other
20 genes within a family of genes exhibiting homology or substantial homology to the target gene.

For antisense suppression, the introduced sequence also need not be full length relative to either the primary transcription product or fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence.
25 Furthermore, the introduced sequence need not have the same intron or exon pattern, and homology of non-coding segments may be equally effective. Normally, a sequence of between about 30 or 40 nucleotides and about full length nucleotides should be used, though a sequence of at least about 100 nucleotides is preferred, a sequence of at least about 200 nucleotides is more preferred, and a sequence of about 500 to about 1700
30 nucleotides is especially preferred.

A number of gene regions can be targetted to suppress *ADC* gene expression. The targets can include, for instance, the coding regions (*e.g.*, regions flanking the PA2 domains), introns, sequences from exon/intron junctions, 5' or 3'

untranslated regions, and the like. In some embodiments, the constructs can be designed to eliminate the ability of regulatory proteins to bind to *ADC* gene sequences that are required for its cell- and/or tissue-specific expression. Such transcriptional regulatory sequences can be located either 5'-, 3'-, or within the coding region of the gene and can be either promote (positive regulatory element) or repress (negative regulatory element) gene transcription. These sequences can be identified using standard deletion analysis, well known to those of skill in the art. Once the sequences are identified, an antisense construct targeting these sequences is introduced into plants to control AP2 gene transcription in particular tissue, for instance, in developing ovules and/or seed.

Oligonucleotide-based triple-helix formation can be used to disrupt *ADC* gene expression. Triplex DNA can inhibit DNA transcription and replication, generate site-specific mutations, cleave DNA, and induce homologous recombination (*see, e.g.,* Havre and Glazer *J. Virology* 67:7324-7331 (1993); Scanlon *et al. FASEB J.* 9:1288-1296 (1995); Giovannangeli *et al. Biochemistry* 35:10539-10548 (1996); Chan and Glazer *J. Mol. Medicine (Berlin)* 75: 267-282 (1997)). Triple helix DNAs can be used to target the same sequences identified for antisense regulation.

Catalytic RNA molecules or ribozymes can also be used to inhibit expression of *ADC* genes. It is possible to design ribozymes that specifically pair with virtually any target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself altered, and is thus capable of recycling and cleaving other molecules, making it a true enzyme. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the constructs. Thus, ribozymes can be used to target the same sequences identified for antisense regulation.

A number of classes of ribozymes have been identified. One class of ribozymes is derived from a number of small circular RNAs which are capable of self-cleavage and replication in plants. The RNAs replicate either alone (viroid RNAs) or with a helper virus (satellite RNAs). Examples include RNAs from avocado sunblotch viroid and the satellite RNAs from tobacco ringspot virus, lucerne transient streak virus, velvet tobacco mottle virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. The design and use of target RNA-specific ribozymes is described in Zhao and Pick *Nature* 365:448-451 (1993); Eastham and Ahlering *J. Urology* 156:1186-1188

(1996); Sokol and Murray *Transgenic Res.* 5:363-371 (1996); Sun *et al. Mol. Biotechnology* 7:241-251 (1997); and Haseloff *et al. Nature*, 334:585-591 (1988).

Another method of suppression is sense cosuppression. Introduction of nucleic acid configured in the sense orientation has been recently shown to be an effective means by which to block the transcription of target genes. For an example of the use of this method to modulate expression of endogenous genes (see, Assaad *et al. Plant Mol. Bio.* 22: 1067-1085 (1993); Flavell *Proc. Natl. Acad. Sci. USA* 91: 3490-3496 (1994); Stam *et al. Annals Bot.* 79: 3-12 (1997); Napoli *et al., The Plant Cell* 2:279-289 (1990); and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184).

The suppressive effect may occur where the introduced sequence contains no coding sequence *per se*, but only intron or untranslated sequences homologous to sequences present in the primary transcript of the endogenous sequence. The introduced sequence generally will be substantially identical to the endogenous sequence intended to be repressed. This minimal identity will typically be greater than about 65%, but a higher identity might exert a more effective repression of expression of the endogenous sequences. Substantially greater identity of more than about 80% is preferred, though about 95% to absolute identity would be most preferred. As with antisense regulation, the effect should apply to any other proteins within a similar family of genes exhibiting homology or substantial homology.

For sense suppression, the introduced sequence, needing less than absolute identity, also need not be full length, relative to either the primary transcription product or fully processed mRNA. This may be preferred to avoid concurrent production of some plants which are overexpressers. A higher identity in a shorter than full length sequence compensates for a longer, less identical sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and identity of non-coding segments will be equally effective. Normally, a sequence of the size ranges noted above for antisense regulation is used. In addition, the same gene regions noted for antisense regulation can be targetted using cosuppression technologies.

Alternatively, *ADC* activity may be modulated by eliminating the proteins that are required for *ADC* cell-specific gene expression. Thus, expression of regulatory proteins and/or the sequences that control *ADC* gene expression can be modulated using the methods described here.

Another method is use of engineered tRNA suppression of *ADC* mRNA translation. This method involves the use of suppressor tRNAs to transactivate target genes containing premature stop codons (*see*, Betzner *et al. Plant J.* 11:587-595 (1997); and Choisne *et al. Plant J.* 11: 597-604 (1997)). A plant line containing a constitutively expressed *ADC* gene that contains an amber stop codon is first created. Multiple lines of plants, each containing tRNA suppressor gene constructs under the direction of cell-type specific promoters are also generated. The tRNA gene construct is then crossed into the *ADC* line to activate *ADC* activity in a targeted manner. These tRNA suppressor lines could also be used to target the expression of any type of gene to the same cell or tissue types.

Some *ADC* proteins (*e.g.*, AP2) are believed to form multimers *in vivo*. As a result, an alternative method for inhibiting *ADC* function is through use of dominant negative mutants. This approach involves transformation of plants with constructs encoding mutant *ADC* polypeptides that form defective multimers with endogenous wild-type *ADC* proteins and thereby inactivate the protein. The mutant polypeptide may vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain. Use of dominant negative mutants to inactivate target genes is described in Mizukami *et al. Plant Cell* 8:831-845 (1996). DNA sequence analysis and DNA binding studies strongly suggests that AP2 (Jofuku *et al.*, *Plant Cell* 6: 1211-1225 (1994)); and several RAP2s function as transcription factors. Thus, dominant-negative forms of *ADC* genes that are defective in their abilities to bind to DNA can also be used.

The AP2 protein is thought to exist in both a phosphorylated and a nonphosphorylated form. Thus AP2 activity may also be regulated by protein kinase signal transduction cascades. In addition, RAP2 gene activity may also be regulated by and/or play a role in protein kinase signal transduction cascades (EREBPs, Ohme-Takagi and Shinshi *Plant Cell* 7: 173-182 (1995); AtEBP, Buttner and Singh *Proc. Natl. Acad. Sci. USA* 94: 5961-5966 (1997); Pti4/5/6, Zhou *et al. EMBO J.* 16: 3207-3218 (1997)). Thus, mutant forms of the *ADC* proteins used in dominant negative strategies can include substitutions at amino acid residues targeted for phosphorylation so as to decrease phosphorylation of the protein. Alternatively, the mutant *ADC* forms can be designed so that they are hyperphosphorylated.

Glycosylation events are known to affect protein activity in a cell- and/or tissue-specific manner (*see*, Meshi and Iwabuchi *Plant Cell Physiol.* 36: 1405-1420 (1995); Meynial-Salles and Combes *J. Biotech.* 46: 1-14 (1996)). Thus, mutant forms of the ADC proteins can also include those in which amino acid residues that are targeted for glycosylation are altered in the same manner as that described for phosphorylation mutants.

AP2 may carry out some of its functions through its interactions with other transcription factors/proteins (*e.g.*, AINTEGUMENTA, Elliott *et al.* *Plant Cell* 8: 155-168 (1996); Klucher *et al.* *Plant Cell* 8: 137-153 (1996); CURLY LEAF, Goodrich *et al.* *Nature (London)* 386: 44-51 (1997); or LEUNIG, Liu and Meyerowitz *Development* 121: 975-991 (1995). Thus, one simple method for suppressing ADC activity is to suppress the activities of proteins that are required for AP2 activity. ADC activity can thus be controlled by "titrating " out transcription factors/proteins required for ADC activity. This can be done by overexpressing domains ADC proteins that are involved in protein:protein interactions in plant cells (*e.g.*, AP2 domains or the putative transcriptional activation domain as described in Jofuku *et al.*, *Plant Cell* 6: 1211-1225 (1994)). This strategy has been used to modulate gene activity (Lee *et al.*, *Exptl. Cell Res.* 234: 270-276 (1997); Thiesen *Gene Expression* 5: 229-243 (1996); and Waterman *et al.*, *Cancer Res.* 56: 158-163 (1996)).

Another strategy to affect the ability of an ADC protein to interact with itself or with other proteins involves the use of antibodies specific to ADC. In this method cell-specific expression of AP2-specific Abs is used inactivate functional domains through antibody:antigen recognition (*see*, Hupp *et al.* *Cell* 83:237-245 (1995)).

Use of nucleic acids of the invention to enhance ADC gene expression

Isolated sequences prepared as described herein can also be used to introduce expression of a particular ADC nucleic acid to enhance or increase endogenous gene expression. Enhanced expression will generally lead to smaller seeds or seedless fruit. Where overexpression of a gene is desired, the desired gene from a different species may be used to decrease potential sense suppression effects.

One of skill will recognize that the polypeptides encoded by the genes of the invention, like other proteins, have different domains which perform different functions. Thus, the gene sequences need not be full length, so long as the desired

functional domain of the protein is expressed. The distinguishing features of *ADC* polypeptides, including the AP2 domain, are discussed in detail below.

Modified protein chains can also be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art and described in detail, below. For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain.

10 Preparation of recombinant vectors

To use isolated sequences in the above techniques, recombinant DNA vectors suitable for transformation of plant cells are prepared. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, for example, Weising et al. *Ann. Rev. Genet.* 15 22:421-477 (1988). A DNA sequence coding for the desired polypeptide, for example a cDNA sequence encoding a full length protein, will preferably be combined with transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the gene in the intended tissues of the transformed plant.

For example, for overexpression, a plant promoter fragment may be employed which will direct expression of the gene in all tissues of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S 25 transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill. Such genes include for example, the *AP2* gene, *ACT11* from *Arabidopsis* (Huang et al. *Plant Mol. Biol.* 33:125-139 (1996)), *Cat3* from *Arabidopsis* (GenBank No. U43147, Zhong et al., *Mol. Gen. Genet.* 251:196-203 30 (1996)), the gene encoding stearyl-acyl carrier protein desaturase from *Brassica napus* (Genbank No. X74782, Solocombe et al. *Plant Physiol.* 104:1167-1176 (1994)), *GPcl* from maize (GenBank No. X15596, Martinez et al. *J. Mol. Biol.* 208:551-565 (1989)),

and *Gpc2* from maize (GenBank No. U45855, Manjunath *et al.*, *Plant Mol. Biol.* 33:97-112 (1997)).

Alternatively, the plant promoter may direct expression of the *ADC* nucleic acid in a specific tissue or may be otherwise under more precise environmental or developmental control. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions, elevated temperature, or the presence of light. Such promoters are referred to here as "inducible" or "tissue-specific" promoters. One of skill will recognize that a tissue-specific promoter may drive expression of operably linked sequences in tissues other than the target tissue. Thus, as used herein a tissue-specific promoter is one that drives expression preferentially in the target tissue, but may also lead to some expression in other tissues as well.

Examples of promoters under developmental control include promoters that initiate transcription only (or primarily only) in certain tissues, such as fruit, seeds, or flowers. Promoters that direct expression of nucleic acids in ovules, flowers or seeds are particularly useful in the present invention. As used herein a seed-specific promoter is one which directs expression in seed tissues, such promoters may be, for example, ovule-specific, embryo-specific, endosperm-specific, integument-specific, seed coat-specific, or some combination thereof. Examples include a promoter from the ovule-specific *BEL1* gene described in Reiser *et al.* *Cell* 83:735-742 (1995) (GenBank No. U39944). Other suitable seed specific promoters are derived from the following genes: *MAC1* from maize (Sheridan *et al.* *Genetics* 142:1009-1020 (1996), *Cat3* from maize (GenBank No. L05934, Abler *et al.* *Plant Mol. Biol.* 22:10131-1038 (1993), the gene encoding oleosin 18kD from maize (GenBank No. J05212, Lee *et al.* *Plant Mol. Biol.* 26:1981-1987 (1994)), *viviparous-1* from *Arabidopsis* (Genbank No. U93215), the gene encoding oleosin from *Arabidopsis* (Genbank No. Z17657), *Atmyc1* from *Arabidopsis* (Urao *et al.* *Plant Mol. Biol.* 32:571-576 (1996), the 2s seed storage protein gene family from *Arabidopsis* (Conceicao *et al.* *Plant* 5:493-505 (1994)) the gene encoding oleosin 20kD from *Brassica napus* (GenBank No. M63985), *napA* from *Brassica napus* (GenBank No. J02798, Josefsson *et al.* *JBL* 26:12196-1301 (1987), the napin gene family from *Brassica napus* (Sjodahl *et al.* *Planta* 197:264-271 (1995), the gene encoding the 2S storage protein from *Brassica napus* (Dasgupta *et al.* *Gene* 133:301-302 (1993)), the genes encoding oleosin A (Genbank No. U09118) and oleosin B (Genbank No. U09119) from

soybean and the gene encoding low molecular weight sulphur rich protein from soybean (Choi *et al. Mol Gen, Genet.* 246:266-268 (1995)).

If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

The vector comprising the sequences (*e.g.*, promoters or coding regions) from genes of the invention will typically comprise a marker gene which confers a selectable phenotype on plant cells. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

Production of transgenic plants

DNA constructs of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment.

Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski *et al. Embo J.* 3:2717-2722 (1984). Electroporation techniques are described in Fromm *et al. Proc. Natl. Acad. Sci. USA* 82:5824 (1985). Ballistic transformation techniques are described in Klein *et al. Nature* 327:70-73 (1987).

Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria. *Agrobacterium tumefaciens*-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, for example Horsch *et al. Science* 233:496-498 (1984), and Fraley *et al. Proc. Natl. Acad. Sci. USA* 80:4803 (1983).

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype such as increased seed mass. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation and Culture, Handbook of Plant Cell Culture*, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee et al. *Ann. Rev. of Plant Phys.* 38:467-486 (1987).

The nucleic acids of the invention can be used to confer desired traits on essentially any plant. Thus, the invention has use over a broad range of plants, including species from the genera *Anacardium*, *Arachis*, *Asparagus*, *Atropa*, *Avena*, *Brassica*, *Citrus*, *Citrullus*, *Capsicum*, *Carthamus*, *Cocos*, *Coffea*, *Cucumis*, *Cucurbita*, *Daucus*, *Elaeis*, *Fragaria*, *Glycine*, *Gossypium*, *Helianthus*, *Heterocallis*, *Hordeum*, *Hyoscyamus*, *Lactuca*, *Linum*, *Lolium*, *Lupinus*, *Lycopersicon*, *Malus*, *Manihot*, *Majorana*, *Medicago*, *Nicotiana*, *Olea*, *Oryza*, *Panieum*, *Pannisetum*, *Persea*, *Phaseolus*, *Pistachia*, *Pisum*, *Pyrus*, *Prunus*, *Raphanus*, *Ricinus*, *Secale*, *Senecio*, *Sinapis*, *Solanum*, *Sorghum*, *Theobromus*, *Trigonella*, *Triticum*, *Vicia*, *Vitis*, *Vigna*, and *Zea*.

Increasing seed size, protein, amino acid, and oils content is particularly desirable in crop plants in which seed are used directly for animal or human consumption or for industrial purposes. Examples include soybean, canola, and grains such as rice, wheat, corn, rye, and the like. Decreasing seed size, or producing seedless varieties, is particularly important in plants grown for their fruit and in which large seeds may be undesirable. Examples include cucumbers, tomatoes, melons, and cherries.

One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

Since transgenic expression of the nucleic acids of the invention leads to phenotypic changes in seeds and fruit, plants comprising the expression cassettes

discussed above must be sexually crossed with a second plant to obtain the final product. The seed of the invention can be derived from a cross between two transgenic plants of the invention, or a cross between a plant of the invention and another plant. The desired effects (*e.g.*, increased seed mass) are generally enhanced when both parental plants
5 contain expression cassettes of the invention.

Seed obtained from plants of the present invention can be analyzed according to well known procedures to identify seed with the desired trait. Increased or decreased size can be determined by weighing seeds or by visual inspection. Protein content is conveniently measured by the method of Bradford *et al. Anal. Bioch.* 72:248
10 (1976). Oil content is determined using standard procedures such as gas chromatography. These procedures can also be used to determine whether the types of fatty acids and other lipids are altered in the plants of the invention.

Using these procedures one of skill can identify the seed of the invention by the presence of the expression cassettes of the invention and increased seed mass.
15 Usually, the seed mass will be at least about 10%, often about 20% greater than the average seed mass of plants of the same variety that lack the expression cassette. The mass can be about 50% greater and preferably at least about 75% to about 100% greater. Increases in other properties *e.g.*, protein and oil will usually be proportional to the increases in mass. Thus, in some embodiments protein or oil content can increase by
20 about 10%, 20%, 50%, 75% or 100%, or in approximate proportion to the increase in mass.

Alternatively, seed of the invention in which *AP2* expression is enhanced will have the expression cassettes of the invention and decreased seed mass. Seed mass will be at least about 20% less than the average seed mass of plants of the same variety
25 that lack the expression cassette. Often the mass will be about 50% less and preferably at least about 75% less or the seed will be absent. As above, decreases in other properties *e.g.*, protein and oil will be proportional to the decreases in mass.

The following Examples are offered by way of illustration, not limitation.

30

Example 1

AP2 Gene Isolation

The isolation and characterization of an *AP2* gene from *Arabidopsis* is described in detail in Jofuku *et al.*, *supra*. Briefly, T-DNA from *Agrobacterium* was

used as an insertional mutagen to identify and isolate genes controlling flower formation in *Arabidopsis*. One transformed line, designated T10, segregated 3 to 1 for a flower mutant that phenotypically resembled many allelic forms of the floral homeotic mutant *ap2*. T10 was tested and it was confirmed genetically that T10 and *ap2* are allelic. The mutant was designated as *ap2-10*.

It was determined that *ap2-10* was the product of a T-DNA insertion mutation by genetic linkage analysis using the T-DNA-encoded neomycin phosphotransferase II (NPTII) gene as a genetic marker. An overlapping set of T-DNA-containing recombinant phage was selected from an *ap2-10* genome library and the plant DNA sequences flanking the T-DNA insertion element were used as hybridization probes to isolate phage containing the corresponding region from a wild-type *Arabidopsis* genome library. The site of T-DNA insertion in *ap2-10* was mapped to a 7.2-kb EcoRI fragment centrally located within the *AP2* gene region.

Five *Arabidopsis* flower cDNA clones corresponding to sequences within the 7.2-kb *AP2* gene region were isolated. All five cloned cDNAs were confirmed to represent *AP2* gene transcripts using an antisense gene strategy to induce *ap2* mutant flowers in wild-type plants.

To determine *AP2* gene structure, the nucleotide sequences of the cDNA inserts were compared to that of the 7.2-kb *AP2* genomic fragment. These results showed that the *AP2* gene is 2.5 kb in length and contains 10 exons and 9 introns that range from 85 to 110 bp in length. The *AP2* gene encodes a theoretical polypeptide of 432 amino acids with a predicted molecular mass of 48 kD. The *AP2* nucleotide and predicted protein sequences were compared with a merged, nonredundant data base. It was found that *AP2* had no significant global similarity to any known regulatory protein.

Sequence analysis, however, did reveal the presence of several sequence features that may be important for *AP2* protein structure or function. First, *AP2* contains a 37-amino acid serine-rich acidic domain (amino acids 14 to 50) that is analogous to regions that function as activation domains in a number of RNA polymerase II transcription factors. Second, *AP2* has a highly basic 10-amino acid domain (amino acids 119 to 128) that includes a putative nuclear localization sequence KKSR suggesting that *AP2* may function in the nucleus. Finally, that the central core of the *AP2* polypeptide (amino acids 129 to 288) contains two copies of a 68-amino acid direct repeat that is referred to here as the *AP2* domain. The two copies of this repeat,

designated *AP2-R1* and *AP2-R2*, share 53% amino acid identity and 69% amino acid homology. Figure 1A shows that each *AP2* repeat contains an 18-amino acid conserved core region that shares 83% amino acid homology. Figure 1B shows that both copies of this core region are theoretically capable of forming amphipathic α -helical structures that may participate in protein-protein interactions. SEQ ID NO:3 is the full length *AP2* genomic sequence.

Example 2

Preparation of *AP2* Constructs

Gene constructs were made comprising the *AP2* gene coding region described above in a transcriptional fusion with the cauliflower mosaic virus 35S constitutive promoter in both the sense and antisense orientations. The original vector containing the 35S promoter pGSJ780A was obtained from Plant Genetic Systems (Gent, Belgium). The pGSJ780A vector was modified by inserting a ClaI-BamHI adaptor containing an EcoRI site in the unique BamHI site of pGSJ780A. The modified pGSJ780A DNA was linearized with EcoRI and the *AP2* gene coding region inserted as a 1.68 kb EcoRI fragment in both sense and antisense orientations with respect to the 35S promoter (*see*, Figures 2 and 3).

The resultant DNA was transformed into *E. coli* and spectinomycin resistant transformants were selected. Plasmid DNAs were isolated from individual transformants and the orientation of the insert DNAs relative to the 35S promoter were confirmed by DNA sequencing. Bacterial cells containing the 35S/*AP2* sense (designated pPW12.4 and pPW9) and 35S/*AP2* antisense (designated pPW14.4 and pPW15) constructs were conjugated to *Agrobacterium tumefaciens* and rifampicin, spectinomycin resistant transformants were selected for use in *Agrobacterium*-mediated plant transformation experiments.

The 35S/*AP2* sense and 35S/*AP2* antisense constructs were introduced into wild-type *Arabidopsis* and tobacco plants according to standard techniques. Stable transgenic plant lines were selected using the plant selectable marker *NPTII* (which confers resistance to the antibiotic kanamycin) present on the modified Ti plasmid vector pGSJ780A.

Example 3

Modification of Seed using AP2 Sequences

This example shows that *ap2* mutant plants and transgenic plants containing the 35S/AP2 antisense construct produced seed with increased mass and total protein content. By contrast, transgenic plants containing the 35S/AP2 *sense* construct produced seed with decreased mass and protein content. Together these results indicate that seed mass and seed contents in transgenic plants can be modified by genetically altering AP2 activity.

Seed from 30 lines were analyzed for altered seed size and seed protein content including the Arabidopsis *ap2* mutants *ap2-1*, *ap2-3*, *ap2-4*, *ap2-5*, *ap2-6*, *ap2-9* and *ap2-10* and transgenic Arabidopsis and transgenic tobacco containing the CaMV 35S/AP2 antisense gene construct, the CaMV 35S/AP2 sense gene construct, or the pGSJ780A vector as described above. The *ap2* mutants used in this study are described in Komaki *et al.*, *Development* 104, 195-203 (1988), Kunst *et al.*, *Plant Cell* 1, 1195-1208 (1989), Bowman *et al.*, *Development* 112, 1-20 (1991), and Jofuku *et al.*, *supra*.

Due to the small size of Arabidopsis and tobacco seed, average seed mass was determined by weighing seed in batches of 100 for Arabidopsis and 50 seed for tobacco. The net change in seed mass due to changes in AP2 gene activity was calculated by subtracting the average mass of wild-type seed from mutant seed mass.

Seed from three wild-type Arabidopsis ecotypes C24, Landsberg-*er*, and Columbia, and one wild-type tobacco SR1 were used as controls. Wild-type Arabidopsis seed display seasonal variations in seed mass which range from 1.6-2.3 mg per 100 seed as shown in Table I. Therefore transgenic Arabidopsis seed were compared to control seed that had been harvested at approximately the same time of season. This proved to be an important for comparing the effects of weak *ap2* mutations on seed mass.

Table I shows that all *ap2* mutant seed examined, *ap2-1*, *ap2-3*, *ap2-4*, *ap2-5*, *ap2-6*, *ap2-9*, and *ap2-10*, show a significant increase in average seed mass ranging from +27 to +104 percent compared to wild-type. The weak partial loss-of-function mutants such as *ap2-1* and *ap2-3* show the smallest gain in average seed mass ranging from +27 percent to +40 percent of wild-type, respectively. By contrast, strong *ap2* mutants such as *ap2-6* and *ap2-10* show the largest gain in seed mass ranging

from +69 percent to +104 percent of wild-type, respectively. Thus reducing *AP2* gene activity genetically consistently increases Arabidopsis seed mass.

AP2 antisense and AP2 sense cosuppression strategies described above were used to reduce *AP2* gene activity *in planta* to determine whether seed mass could be manipulated in transgenic wild-type plants. Twenty-nine independent lines of transgenic Arabidopsis containing the CaMV 35S/AP2 antisense gene constructs pPW14.4 and pPW15 (Figure 2) were generated. Each transgenic line used in this study tested positive for kanamycin resistance and the presence of one or more copies of T-DNA.

Table I shows that seed from nine transgenic Arabidopsis AP2 antisense lines show a significant increase in seed mass when compared to control seed ranging from +22 percent for line C24 15-542 to +89 percent for line C24 15-566. Both C24 and Landsberg-*er* ecotypes were used successfully. Increased seed mass was observed in F1, F2, and F3 generation seed.

Eight lines containing the 35S/AP2 sense gene construct were generated which were phenotypically cosuppression mutants. As shown in Table I seed from two cosuppression lines examined showed larger seed that range from +26 percent to +86 percent. By contrast, plants transformed with the vector pGSJ780A showed a normal range of average seed mass ranging from -0.5 percent to +13 percent compared to wild-type seed (Table I). Together, these results demonstrate that *AP2* gene sequences can be used to produce a significant increase in Arabidopsis seed mass using both antisense and cosuppression strategies in a flowering plant.

Table I. Genetic control of Arabidopsis seed mass by AP2.

		Average seed mass in mg per 100 seed ^{1,2}	Percent change in seed mass compared to wild-type
5	ap2 mutant seed	2.1 (0.1)	+27%
	1. <i>ap2-1</i>	2.2 (0.1)	+33%
		2.1 (0.2)	+31%
		2.8 (0.2)	+33%
10	2. <i>ap2-3</i>	2.6 (0.1)	+27%
	3. <i>ap2-4</i>	3.5 (0.3)	+69%
		3.5 (0.2)	+69%
	4. <i>ap2-5</i>	2.9 (0.1)	+39%
	5. <i>ap2-6</i>	3.5 (0.2)	+69%
15	6. <i>ap2-9</i>	2.9 (0.1)	+40%
	7. <i>ap2-10</i>	3.7 (0.4)	+79%
		3.9 (0.3)	+90%
		4.2 (0.5)	+104%
Seed produced by transgenic CaMV35S/AP2 antisense lines (from a Km resistant mother)			
20	1. C24 14.4E (F1-15) F2 sd		+35%
	C24 14.4E (F1-15) F3 sd	3.1	+47%
	2. C24 14.4S (F1-1)	3.4 (0.3)	+29%
	3. C24 14.4AA (F1-24)	2.8 (0.2)	+30%
	4. C24 14.4DD (F1-2)	2.9 (0.1)	+30%
25	5. C24 15-522	2.8 (0.3)	+76%
	6. C24 15-542 (F1-2)	3.6 (0.1)	+25%
	C24 15-542 (F1-7)	2.6 (0.1)	+22%
	7. C24 15-566	2.5 (0.2)	+89%
	8. LE 15-9992-3 (F1-1) F2 sd	3.9 (0.1)	+42%
30	9. LE 15-83192-3 (F1-3)	2.4 (0.1)	+33%
	LE 15-83192-3 (F1-17)	2.8 (0.0)	+28%
		2.7 (0.0)	
Seed produced by transgenic CaMV35S/AP2 cosuppression lines (from a Km resistant mother)			
35	1. C24 9-5 (F1-5)	3.8 (0.0)	+86%
	2. LE 9-83192-2 (F1-19)	2.7 (0.2)	+26%
	LE 9-83192-2 (F1-24)	2.7 (0.1)	+26%

		Average seed mass in mg per 100 seed ^{1,2}	Percent change in seed mass compared to wild-type
Seed produced by transgenic pGSJ780A vector only lines (from a Km resistant mother plant)			
5	1. C24 3 -107 (F1-1)	2.2 (0.1)	+9%
	2. C24 3-109 (F1-1)	2.3 (0.0)	+13%
	3. LE 3-83192-1 (F1-2)	2.3 (0.1)	+7%
	4. LE 3-83192-3 (F1-2)	2.4 (0.1)	+11%
10	5. LE 3-9992-4 (F1-4)	2.4 (0.2)	+12%
	LE 3-9992-4 (F1-6)	2.3 (0.0)	+9%
	LE 3-9992-4 (F1-8)	2.1 (0.0)	-0.5%
	6. LE 3-9992-9 (F1-3)	2.3 (0.1)	+7%
15	Seed produced by wild-type Arabidopsis plants		
	1. C24	2.0 (0.1)	
		2.3 (0.1)	
		2.2	
	2. Landsberg- <i>er</i>	1.6 (0.1)	
20		2.1 (0.1)	
		2.1	
		2.3 (0.1)	
25	3. Columbia	1.8 (0.1)	
		2.1 (0.1)	

¹ Standard deviation values are given in parentheses.

² Wild-type seed values used for this comparison were chosen by ecotype and harvest date.

Arabidopsis *AP2* gene sequences were also used to negatively control seed mass in tobacco, a heterologous plant species. Table II shows that in five transgenic tobacco lines the CaMV 35S/*AP2* overexpression gene construct was effective in reducing transgenic seed mass from -27 percent to -38 percent compared to wild-type seed. These results demonstrate the evolutionary conservation of *AP2* gene function at the protein level for controlling seed mass in a heterologous system.

Table II. Genetic control of tobacco seed mass using Arabidopsis AP2.

	Average seed mass in mg per 5 seed ¹	Percent change in seed mass compared to wild-type
5	Seed produced by transgenic CaMV 35S/AP2 sense gene lines (from a Km resistant mother)	
	1. SR1 9-110 To	3.1 (0.0) -27%
	SR1 9-110 (F1-5)	3.0 (0.2) -29%
10		2.8 (0.3) -34%
	2. SR1 9-202 (F1-G)	3.1 (0.2) -27%
	SR1 9-202 (F1-I)	3.2 (0.1) -24%
	3. SR1 9-103 (F1-2)	3.9 (0.0) -8%
	4. SR1 9-413-1	2.8 (0.0) -34%
15		3.0 (0.2) -29%
	5. SR1 9-418-1 To	3.5 (0.1) -18%
	Seed produced by transgenic CaMV 35S/AP2 antisense gene lines (from a Km resistant mother)	
20	1. SR1 15-111	5.1 (0.4) +20%
	SR1 15-111 (F1)	5.0 (0.4) +19%
	2. SR1 15-116 To	4.1 (0.4) -3%
	SR1 15-116 (F1-2)	4.0 (0.1) -5%
	SR1 15-116 (F1-1)	4.5 (0.1) +5%
25	3. SR1 15-407 (F1)	4.8 (0.5) +10%
		4.7 (0.3) +10%
	4. SR1 15-102 (F1)	4.5 (0.2) +6%
	5. SR1 15-413 (F1-3)	4.2 (0.0) +0%
	6. SR1 15-410 (F1-2)	4.4 (0.0) +4%
30		
	7. SR1 15-210 (F1-4)	3.6 (0.1) -15%
	Seed produced by pGSJ780A vector only lines (from a Km resistant mother)	
35	1. SR1 3-402 (F1)	5.0 (0.1) +17%
	2. SR1 3-401 (F1)	4.6 (0.1) +8%
	3. SR1 3-405 (F1)	4.4 (0.1) +4%
40	Seed from wild-type tobacco	
	1. SR1	4.2 (0.3)
		4.0 (0.1)

¹ Standard deviation values are given in parentheses.

45

Use of AP2 gene constructs to control seed protein content

Total seed protein was extracted and quantitated from seed produced by wild-type, ap2 mutant, transgenic AP2 antisense, and transgenic AP2 sense cosuppression plants according to Naito *et al. Plant Mol Biol.* 11, 109-123 (1988). Seed protein was extracted in triplicate from batches of 100 dried seed for Arabidopsis or 50 dried seed for tobacco. Total protein yield was determined by the Bradford dye-binding procedure as described by Bradford, *Anal. Biochem.* 72:248 (1976). The results of this analysis are shown in Table III.

ap2 mutant total seed protein content increased by 20 percent to 78 percent compared to wild-type control seed. Total seed protein from transgenic *AP2* antisense plants increased by +31 percent to +97 percent compared to wild-type controls. Transgenic *AP2* cosuppression seed showed a +13 and +17 percent increase over wild-type. Together, the transgenic antisense and cosuppression mutant seed consistently yielded more protein per seed than did the wild-type controls or transgenic plants containing the pGSJ780A vector only (Table III).

Table III. Genetic control of total seed protein content in Arabidopsis using *AP2*.

	Total seed protein in μ g per 100 seed ¹	Percent change in protein content compared to wild-type
<i>ap2</i> mutant seed		
1. <i>ap2-1</i>	652 (17)	+20% relative to WT seed
	615 (30)	+11%
2. <i>ap2-3</i>	705 (47)	+27%
3. <i>ap2-4</i>	729 (107)	+33%
4. <i>ap2-5</i>	617 (24)	+13%
5. <i>ap2-6</i>	836 (14)	+52%
6. <i>ap2-9</i>	798 (11)	+46%
7. <i>ap2-10</i>	836 (15)	+78%
Transgenic CaMV 35S/ <i>AP2</i> antisense seed mass (from Km resistant mother)		
1. C24 14.4E (F1-1) F3 sd	615 (60)	+31%
2. C24 15-522 (F1-1)	790 (23)	+68%
3. C24 15-566	925 (173)	+97%
Transgenic CaMV 35S/ <i>AP2</i> sense cosuppression seed mass (from Km Resistant mother plant)		
1. LE 9-83192-2 (F1-19)	616	+13%
LE 9-83192-2 (F1-24)	637	+17%
Wild-type seed		
1. C24	469 (19)	
2. LE	545 (22)	
	555	
3. Col	548 (42)	

¹ Standard deviation values are given in parentheses.

Transgenic tobacco containing the 35S/*AP2* sense gene construct show that *AP2* overexpression can decrease seed protein content by 27 to 45 percent compared to wild-type seed. Together, the transgenic Arabidopsis and tobacco results demonstrate that seed mass and seed protein production can be controlled by regulating *AP2* gene activity.

Table IV. Negative control of transgenic tobacco seed protein content by Arabidopsis AP2 gene expression.¹

	Ave. protein per 50 seed	Percent change in protein content compared to wild-type
Seed produced by transgenic CAMV 35S/AP2 sense gene plant		
1. SR1 9-110	242 (11)	-45%
2. SR1 9-202 (F1-G)	271 (11)	-38%
3. SR1 9-413	362 (8)	-18%
4. SR1 9-418-1	319 (16)	-27%
Wild-type Control SR1 (wild-type)	440 (8) (JO)	NA

¹ Standard deviation values are given in parentheses.

Analysis of transgenic seed proteins by gel electrophoresis

Arabidopsis seed produce two major classes of seed storage proteins, the 12S cruciferins and 2S napins which are structurally related to the major storage proteins found in the *Brassicaceae* and in the *Leguminosae*. The composition of seed proteins in wild-type, *ap2* mutant, and transgenic Arabidopsis seed were compared by SDS polyacrylamide gel electrophoresis as described by Naito *et al.*, *Plant Mol. Biol.* 11, 109-123 (1988). Total seed proteins were extracted as described above. 50 µg aliquots were separated by gel electrophoresis and stained using Coomassie brilliant blue. These results showed that the spectrum of proteins in wild-type and *ap2* mutant seed are qualitatively indistinguishable. There is no detectable difference in the representation of the 12S or 2S storage proteins between the wild-type and *ap2* mutant seed extracts. This shows that reducing *AP2* gene activity genetically does not alter the profile of storage proteins synthesized during seed maturation. The spectrum of seed proteins produced in transgenic *AP2* antisense and *AP2* sense cosuppression seed are also indistinguishable from wild-type. In particular, there is no detectable difference in the representation of the 12S cruciferin or 2S napin storage proteins in the larger seed.

Finally, the transgenic tobacco plants containing the 35S/*AP2* overexpression gene construct produced significantly smaller seed. Despite the decrease in seed mass in transgenic tobacco there was no detectable difference in storage protein profiles between seed from 35S/*AP2* transformants and wild-type SR1.

Example 4

Isolation of other members of the AP2 gene family from *Arabidopsis*

This example describes isolation of a number of *AP2* nucleic acids from *Arabidopsis*. The nucleic acids are referred to here as *RAP2* (related to *AP2*) were identified using primers specific to nucleic acid sequences from the *AP2* domain described above.

MATERIALS AND METHODS

Plant Material. *Arabidopsis thaliana* ecotype Landsberg *erecta* (L-er) and C24 were used as wild type. Plants were grown at 22°C under a 16-hr light/8-hr dark photoperiod in a 1:1:1 mixture containing vermiculite/perlite/peat moss. Plants were watered with a one-fourth strength Peter's solution (Grace-Sierra, Milpitas, CA). Root tissue was harvested from plants grown hydroponically in sterile flasks containing 1x Murashige and Skoog plant salts (GIBCO), 1 mg/liter thiamine, 0.5 mg/liter pyridoxine, 0.5 mg/liter nicotinic acid, 0.5 g/liter 2-(*N*-morpholino)ethanesulfonic acid (MES), and 3% sucrose, with moderate shaking and 70 $\mu\text{mol} \cdot \text{m}^{-2} \cdot \text{sec}^{-1}$ of light.

Analysis of cloned *Arabidopsis* cDNAs. *Arabidopsis* expressed sequence tagged (EST) cDNA clones representing *RAP2.1* and *RAP2.9* were generated as described by Cooke *et al.* (Cooke, R., *et al.*, 1996, *Plant J.* 9, 101-124). EST cDNA clones representing *RAP2.2* and *RAP2.8* were generated as described by Höfte *et al.* (Höfte, H., *et al.*, 1993, *Plant J.* 4, 1051-1061). EST cDNA clones representing all other *RAP2* genes were generated by Newman *et al.* (Newman, T., *et al.*, 1994, *Plant Physiol.* 106,1241-1255) and provided by the *Arabidopsis* Biological Resource Center (Ohio State University). Plasmid DNAs were isolated and purified by anion exchange chromatography (Qiagen, Chatsworth, CA). DNA sequences were generated using fluorescence dye-based nucleotide terminators and analyzed as specified by the manufacturer (Applied Biosystems).

Nucleotide and Amino Acid Sequence Comparisons. The TBLASTN program (Altschul, S. F., *et al.*, 1990, *J. Mol. Biol.* 215, 403-410) and default parameter settings were used to search the *Arabidopsis* EST database (AAtdB 4-7) for genes that encode *AP2* domain-containing proteins. Amino acid sequence alignments were generated using the CLUSTAL W multiple sequence alignment program (Thompson, J. D., *et al.*, 1994, *Nucleic Acids Res.* 22, 4673-4680). Secondary structure

predictions were based on the principles and software programs described by Rost (Rost, B., 1996, *Methods Enzymol.* 266, 525-539) and Rost and Sander (Rost, B., *et al.*, 1993, *J. Mol. Biol.* 232, 589-599; Rost, B., *et al.*, 1994, *Proteins* 19, 55-77).

RAP2 Gene-Specific Probes. RAP2 gene-specific fragments were

5 generated by PCR using gene-specific primers and individual RAP2 plasmid DNAs as a template as specified by Perkin-Elmer (Roche Molecular Systems, Branchburg, NJ). The following primers were used to generate fragments representing each RAP2 gene:

RAP2.1, 5'-AAGAGGACCATCTCTCAG-3',

5'-AACACTCGCTAGCTTCTC-3';

10 RAP2.2, 5'-TGGTTCAGCAGCCAACAC-3', 5'-CAATGCATAGAGCTTGAGG-3';

RAP2.3, 5'-TCATCGCCACGATCAACC-3', 5'-AGCAGTCCAATGCGACGG-3';

RAP2.4, 5'-ACGGATTTCACATCGGAG-3', 5'-CTAAGCTAGAATCGAATCC-3';

RAP2.7, 5'-CGATGGAGACGAAGACTC-3, 5'-GTCGGAACCGGAGTTACC-3';

RAP2.8, 5'-TCACTCAAAGGCCGAGATC-3', 5'-TAACAACATCACCGGCTCG-3';

15 RAP2.9, 5'-GTGAAGGCTTAGGAGGAG-3', 5'-TGCCTCATATGAGTCAGAG-3'.

PCR-synthesized DNA fragments were gel purified and radioactively labeled using random oligonucleotides (Amersham) for use as probes in gene mapping and RNA gel blot experiments.

Gene Mapping Experiments. RAP2 genes were placed on the

20 *Arabidopsis* genetic map by either restriction fragment length polymorphism segregation analysis using recombinant inbred lines as described by Reiter *et al.* (Reiter, R. S., *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89, 1477-1481) or by matrix-based analysis of pooled DNAs from the *Arabidopsis* yUP or CIC yeast artificial chromosome (YAC) genomic libraries (Ecker, J. R., 1990, *Methods* 1, 186-194; Creusot, F., *et al.*, 1995, 25 *Plant J.* 8, 763-770) using the PCR (Green, E. D., *et al.*, 1990, *Proc. Natl. Acad. Sci. USA* 87, 1213-1217; Kwiatkowski, T. J., *et al.*, 1990, *Nucleic Acids Res.* 18, 7191-7192). Matrix based mapping results were confirmed by PCR using DNA from individual YAC clones.

mRNA Isolation. Polysomal poly(A) mRNAs from *Arabidopsis* flower,

30 rosette leaf, inflorescence stem internode, and hydroponically-grown roots were isolated according to Cox and Goldberg (Cox, K. H., *et al.*, 1988, in *Plant Molecular Biology. A Practical Approach*, ed. Shaw, C. H. (IRL, Oxford), pp. 1-35).

RNA Gel Blot Studies. RNA gel blot hybridizations were carried out as specified by the manufacturer (Amersham). mRNA sizes were estimated relative to known RNA standards (BRL). AP2 transcripts were detected using a labeled DNA fragment representing nucleotides 1-1371 of the AP2 cDNA plasmid clone pAP2cl (Jofuku, K. D., *et al.*, 1994, *Plant Cell* 6, 1211-1225).

RESULTS

The AP2 Domain Defines a Large Family of Plant Proteins. Using the AP2 domain as a sequence probe 34 cDNA clones were identified that encode putative RAP2 proteins in the Arabidopsis EST database (*Materials and Methods*). Several of these partial sequences have been reported previously (Ohme-Takagi, *et al.*, 1995, *Plant Cell* 7, 173-182; Elliot, R. C., *et al.*, 1996, *Plant Cell* 8, 155-168; Klucher, K M., *et al.*, 1996, *Plant Cell* 8, 137-153; Wilson, K., *et al.*, 1996, *Plant Cell* 8, 659-671; Ecker, J. R., 1995, *Science* 268, 667-675; Weigel, D., 1995, *Plant Cell* 7, 388-389). Based on nucleotide sequence comparison, it was inferred that approximately half of the 34 RAP2 cDNA sequences were likely to represent redundant clones. Therefore, a complete DNA sequence for 17 putative RAP2 cDNA clones that appeared to represent unique genes and which contained the largest cDNA inserts was selected and generated. It was determined from the predicted amino acid sequences of these clones that the *Arabidopsis* RAP2 ESTs represent a minimum of 12 genes that are designated *RAP2.1-RAP2.12*. As shown in Table V, preliminary gene mapping experiments using restriction fragment length polymorphism analysis and PCR-based screening of the *Arabidopsis* yUP and CIC yeast artificial chromosome libraries (*Materials and Methods*) revealed that at least 7 members of the RAP2 gene family are distributed over 4 different chromosomes. In addition, several family members are tightly linked in the genome. For example, *RAP2.10* is only 10 kb away from *AP2*, which is also closely linked to *ANT* on chromosome 4 (Elliot, R. C., *et al.*, 1996, *Plant Cell* 8, 155-168; Klucher, K M., *et al.*, 1996, *Plant Cell* 8, 137-153).

Sequence analysis also revealed that the proteins encoded by the RAP2 genes are all characterized by the presence of least one AP2 domain. Fig. 4 shows a sequence comparison of 21 AP2 domains from 19 different polypeptides including *RAP2.1-RAP2.12*, *AP2*, *ANT*, *TINY*, and the tobacco EREBPs. From this comparison, it was determined that there are 2 conserved sequence blocks within each AP2 domain.

The first block, referred to as the YRG element, consists of 19-22 amino acids, is highly basic and contains the conserved YRG amino acid motif (Fig. 4 A and B). The second block, referred to as the RAYD element, is 42-43 amino acids in length and contains a highly conserved 18-amino acid core region that is predicted to form an amphipathic α -helix in the AP2 domains of AP2, ANT, TINY, and the EREBPs. In addition, there are several invariant amino acid residues within the YRG and RAYD elements that may also play a critical role in the structure or function of these proteins. For example, the glycine residue at position 40 within the RAYD element is invariant in all AP2 domain containing proteins (Fig. 4 A and B) and has been shown to be important for AP2 function (Jofuku, K. D., *et al.*, 1994, *Plant Cell* 6, 1211-1225).

Table V. *Arabidopsis* RAP2 genes

15	Gene	RAP2 gene containing YAC clones*	Chromosome map position†
20	AINTÉGUMENTA	ND	4-73
	TINY	ND	5-32 to 5-45
25	RAP2.1	yUP18H2, CIC11D10	ND‡
	RAP2.2	yUP6C1 yUP12G6, yUP24B8, yUP23E11, CIC4H5,	38§
	RAP2.3	CIC12C2	3-21
30	RAP2.4	CIC7D2, CIC10C4	ND‡
	RAP2.7	yUP10E1	ND‡
35	RAP2.8	CIC10G7	1-94 to 1-103§
	RAP2.9	CIC9E12	1-117§
	RAP2.10¶	ND	4-73

* YAC clones were determined to contain the specified RAP2 gene by PCR-based DNA synthesis using gene-specific primers (Green, E. D., *et al.*, 1990, *Proc. Natl. Acad. Sci. USA* 87, 1213-1217; Kwiatkowski, T. J., *et al.*, 1990, *Nucleic Acids Res.* 18, 7191-7192).

† Chromosome map positions are given with reference to the *Arabidopsis* unified genetic map (AAtdB 4-7).

‡ YAC-based map position is ambiguous.

§ Preliminary map position is based on a single contact with the physical map.

GenBank accession numbers for complete EST sequences for RAP2 and other genes are as follows: *AINTÉGUMENTA* (U40256/U41339); *TINY*, (X94598), *RAP2.1*

(AF003094), RAP2.2 (AF003095), RAP2.3 (AF003096), RAP2.4 (AF003097), RAP2.5 (AF003098), RAP2.6 (AF003099), RAP2.7 (AF003100), RAP2.8 (AF003101), RAP2.9 (AF003102), RAP2.10 (AF003103), RAP2.11 (AF003104), and RAP2.12 (AF003105).

All RAP2 cDNA clones were originally reported with partial sequences and given

5 GenBank accession numbers as shown in parentheses following each gene name: *RAP2.1* (Z27045), *RAP2.2* (Z26440). *RAP2.3* (T04320 and T13104), *RAP2.4* (T13774), *RAP2.5* (T45365), *RAP2.6* (T45770), *RAP2.7* (T20443), *RAP2.8* (Z33865), *RAP2.9* (Z37270), *RAP2.10* (T76017), *RAP2.11* (T42962), and *RAP2.12* (T42544). Due to the preliminary nature of the EST sequence data, the predicted amino acid sequences for EST Z27045,
10 T04320, T13774, and T42544 contained several errors and were incorrectly reported (Ohme-Takagi, *et al.*, 1995, *Plant Cell* 7, 173-182; Klucher, K M., *et al.*, 1996, *Plant Cell* 8, 137-153; Wilson, K., *et al.*, 1996, *Plant Cell* 8, 659-671; Ecker, J. R., 1995, *Science* 268, 667-675; Weigel, D., 1995, *Plant Cell* 7, 388-389). They are correctly given in the GenBank accession numbers noted above.

15 RAP2 cDNA sequence comparison also shows that there are at least two branches to the RAP2 gene family tree. The AP2-like and EREBP-like branches are distinguished by the number of AP2 domains contained within each polypeptide and by sequences within the conserved YRG element. The AP2-like branch of the RAP2 gene family is comprised of three genes *AP2*, *ANT*, and *RAP2.7*, each of which encodes a
20 protein containing two AP2 domains (Fig. 4A). In addition, these proteins possess a conserved WEAR/WESH amino acid sequence motif located in the YRG element of both AP2 domain repeats (Fig. 4A). By contrast, genes belonging to the EREBP-like branch of the RAP2 gene family encode proteins with only one AP2 domain and include *RAP2.1-RAP2.6*, *RAP2.8-RAP2.12*, and *TINY* (Fig. 4B). Proteins in this class possess a
25 conserved 7-amino acid sequence motif referred to as the WAAEIRD box (Fig. 4B) in place of the WEAR/WESH motif located in the YRG element (Fig. 4A). Based on these comparisons, separate AP2 domain consensus sequences for both classes of RAP2 proteins were generated (Fig. 4 A and B). These results suggest that the AP2 domain and specific sequence elements within the AP2 domain are important for RAP2 protein
30 functions.

The AP2-like class of RAP2 proteins is also characterized by the presence of a highly conserved 25-26 amino acid linker region that lies between the two AP2 domain repeats (Klucher, K M., *et al.*, 1996, *Plant Cell* 8, 137-153). This region is 40%

identical and 48% similar between AP2, ANT and RAP2.7 and is not found in proteins belonging to the EREBP-like branch of RAP2 proteins. Molecular analysis of the *ant-3* mutant allele showed that the invariant C-terminal glycine residue within this linker region is essential for ANT function *in vivo* (Klucher, K M., *et al.*, 1996, *Plant Cell* 8, 137-153), suggesting that the linker region may also play an important role in AP2 and RAP2.7 function.

Sequences Within the RAYD Element are Predicted to Form

Amphipathic α -Helices. As noted above, the 18-amino acid core region within the RAYD element of the AP2 domain in AP2 is predicted to form an amphipathic α -helix that may be important for AP2 structure or function. Secondary structure prediction analysis was used to determine whether this structure has been conserved in RAP2 proteins. As shown in Fig. 4, the core region represents the most highly conserved sequence block in the RAYD element of AP2 and the RAP2 proteins. Secondary structure analysis predicts that all RAP2 proteins contain sequences within the RAYD element that are predicted to form amphipathic α -helices (Fig. 4 A and B). Fig. 4C shows that sequences in RAP2.7-R1 are predicted to form an amphipathic α -helix that is 100% identical to that predicted for AP2-R1 and 63% similar to that predicted for ANT-R1. Sequences within the AP2 domain of EREBP-like RAP2 proteins are predicted to form similar α -helical structures. Fig. 4D shows that the RAP2.2, RAP2.5, and RAP2.12 α -helices are 81, 100, and 81% similar to that predicted for EREBP-3, respectively. Together, these results strongly suggest that the predicted amphipathic α -helix in the RAYD element is a conserved structural motif that is important for AP2 domain function in all RAP2 proteins.

RAP2 Genes are Expressed in Floral and Vegetative Tissues. Previous studies have shown that AP2 and ANT are differentially expressed at the RNA level during plant development (Jofuku, K. D., *et al.*, 1994, *Plant Cell* 6, 1211-1225; Elliot, R. C., *et al.*, 1996, *Plant Cell* 8, 155-168; Klucher, K M., *et al.*, 1996, *Plant Cell* 8, 137-153). AP2 is expressed at different levels in developing flowers, leaves, inflorescence stems, and roots. To determine where in plant development the EREBP-like class of RAP2 genes are expressed RAP2.1, RAP2.2, RAP2.3, and RAP2.4 gene-specific probes were reacted with a mRNA gel blot containing flower, leaf, inflorescence stem, and root polysomal poly(A) mRNA. Results from these experiments showed that each RAP2 gene produces a uniquely sized mRNA transcript and displays a

distinct pattern of gene expression in flowers, leaves, inflorescence stems, and roots. For example, the *RAP2.1* gene is expressed at low levels in wild-type flower, leaf, stem, and root. *RAP2.2* gene expression appears to be constitutive in that *RAP2.2* transcripts are detected at similar levels in wild-type flower, leaf, stem, and root. By contrast, the *RAP2.3* gene is expressed at a low level in wild-type flowers, at a slightly higher level in leaves, and is relatively highly expressed in both stems and roots. Finally, the *RAP2.4* gene is also expressed in wild-type flower, leaf, stem, and root and is most highly expressed in roots and leaves. These data indicate that individual members of the EREBP-like family of RAP2 genes are expressed at the mRNA level in both floral and vegetative tissues and show quantitatively different patterns of gene regulation.

RAP2 Gene Expression Patterns are Affected by *ap2*. RAP2 gene expression was analyzed in *ap2-10* mutant plants by RNA gel blot analysis to determine whether AP2 is required for RAP2 gene expression. The expression of three RAP2 genes are differentially affected by the loss of AP2 function. For example, *RAP2.2* gene expression is not dramatically altered in mutant flowers, leaves, and roots compared to wild-type Landsberg *erecta* but is down-regulated in mutant stem. *RAP2.3* gene expression appears unchanged in mutant roots but is up-regulated in mutant flowers and leaves and down-regulated in mutant stems. By contrast, *RAP2.4* gene expression appears relatively unchanged in mutant stems and roots but is slightly up-regulated in mutant flowers and leaves. To control for possible secondary effects of ecotype on RAP2 gene expression, RAP2 gene expression levels in wild-type C24 and *ap2-10* mutant stems were compared. These results show that the differences in *RAP2.2*, *RAP2.3*, and *RAP2.4* gene expression in C24 and *ap2-10* stem are similar to those observed between wild-type Landsberg *erecta* and *ap2-10* mutant stem. Together these results suggest that AP2 directly or indirectly regulates the expression of at least three RAP2 genes. More importantly, these results suggest that AP2 is controlling gene expression during both reproductive and vegetative development.

DISCUSSION

RAP2 Genes Encode a New Family of Putative DNA Binding Proteins.

One important conclusion from the characterization of these clones is that the AP2 domain has been evolutionarily conserved in at least *Arabidopsis* and tobacco. In addition, there are two subfamilies of AP2 domain containing proteins in *Arabidopsis* that

are designated as the AP2-like and the EREBP-like class of RAP2 proteins. *In vitro* studies have shown that both the EREBP and the AP2 proteins bind to DNA in a sequence specific manner and that the AP2 domain is sufficient to confer EREBP DNA binding activity (Ohme-Takagi, *et al.*, 1995, *Plant Cell* 7, 173-182). From these results and the high degree of sequence similarity between the AP2 domain motifs in AP2, the EREBPs, and the RAP2 proteins, it is concluded that RAP2 proteins function as plant sequence specific DNA binding proteins. Although the exact amino acid residues within the AP2 domain required for DNA binding have not yet been identified, sequence comparisons have revealed two highly conserved motifs referred to as the YRG and RAYD elements within the AP2 domain.

The RAYD element is found in all known AP2 domains and contains a conserved core region that is predicted to form an amphipathic α -helix (Fig. 4). One hypothesis for the function of this α -helical structure is that it is involved in DNA binding, perhaps through the interaction of its hydrophobic face with the major groove of DNA (Zubay, G., *et al.*, 1959, *J. Mol. Biol.* 7, 1-20). Alternatively, this structure may mediate protein-protein interactions important for RAP2 functions. These interactions may involve the ability to form homo- or heterodimers similar to that observed for the MADS box family of plant regulatory proteins (Huang, H., *et al.*, 1996, *Plant Cell* 8, 81-94; Riechmann, J. L., *et al.*, 1996, *Proc. Natl. Acad. Sci. USA* 93, 4793-4798) and for the mammalian ATF/CREB family of transcription factors (Hai, T., *et al.*, 1991, *Proc. Natl. Acad. Sci. USA* 88, 3720-3724; O'Shea, E. K., *et al.*, 1992, *Cell* 68, 699-708.).

The conserved YRG element may also function in DNA binding due to the highly basic nature of this region in all RAP2 proteins (Fig. 4). However, the YRG element also contains sequences that are specific for each class of RAP2 protein and may be functionally important for DNA binding. Specifically, the WAAIERD motif is highly conserved in tobacco EREBPs and in EREBP-like RAP2 proteins. By contrast, the WEAR/WESH motif replaces the WAAIERD box in AP2-like RAP2 proteins (Fig.4). *In vitro* studies suggest that the EREBPs and AP2 recognize distinct DNA sequence elements (Ohme-Takagi, *et al.*, 1995, *Plant Cell* 7, 173-182). It is possible that the WAAIERD and WEAR/WESH motifs may be responsible for DNA binding sequence specificity. The presence of two AP2 domains in AP2 may also contribute to differences in sequence specificity. Although the molecular significance of having one or two AP2

domain motifs is not yet known, genetic and Molecular studies have shown that mutations in either AP2 domain affect AP2 function, implying that both are required for wild-type AP2 activity (Jofuku, K. D., *et al.*, 1994, *Plant Cell* 6, 1211-1225).

In addition to *Arabidopsis* and tobacco, cDNAs that encode diverse AP2 domain-containing proteins have been found in maize, rice, castor bean, and several members of the Brassicaceae including canola (Ohme-Takagi, *et al.*, 1995, *Plant Cell* 7, 173-182; Elliot, R. C., *et al.*, 1996, *Plant Cell* 8, 155-168; Klucher, K M., *et al.*, 1996, *Plant Cell* 8, 137-153; Wilson, K., *et al.*, 1996, *Plant Cell* 8, 659-671 and Weigel, D., 1995, *Plant Cell* 7, 388-389). This strongly suggests that the AP2 domain is an important and evolutionarily conserved element necessary for the structure or function of these proteins.

RAP2 Gene Expression in Floral and Vegetative Tissues. The *AP2*, *RAP2.1*, *RAP2.2*, *RAP2.3*, and *RAP2.4* genes show overlapping patterns of gene expression at the mRNA level in flowers, leaves, inflorescence stems, and roots. However, each gene appears to be differentially regulated in terms of its mRNA prevalence. The overlap in *RAP2* gene activity could affect the genetic analysis of *AP2* and *RAP2* gene functions if these genes are also functionally redundant. For example, in flower development *AP2* and *ANT* show partially overlapping patterns of gene expression at the organ and tissue levels (Jofuku, K. D., *et al.*, 1994, *Plant Cell* 6, 1211-1225; Elliot, R. C., *et al.*, 1996, *Plant Cell* 8, 155-168; Klucher, K M., *et al.*, 1996, *Plant Cell* 8, 137-153; W. Szeto). From single and double mutant analysis it has also been suggested that *AP2* may be partially redundant in function with *ANT* (Elliot, R. C., *et al.*, 1996, *Plant Cell* 8, 155-168). The phenomenon of genetic redundancy and its ability to mask the effects of gene mutation is more clearly demonstrated by the MADS domain containing floral regulatory genes *APETALA1* (*API*) and *CAULIFLOWER* (*CAL*). Genetic studies have demonstrated that mutations in *cal* show no visible floral phenotype except when in double mutant combination with *ap1* (Bowman, J. L., *et al.*, 1993, *Development Cambridge, U.K.*, 119, 721-743), indicating that *API* is completely redundant in function for *CAL*. The hypothesis that the *RAP2* genes may have genetically redundant functions is supported by the fact that the dominant gain-of-function mutation *tiny* is the only *Arabidopsis* *RAP2* EREBP-like gene mutant isolated to date (Wilson, K., *et al.*, 1996, *Plant Cell* 8, 659-671).

AP2 Activity Is Detectable in Vegetative Development. The present analysis of RAP2 gene expression in wild-type and *ap2-10* plants suggests that AP2 contributes to the regulation of RAP2 gene activity throughout *Arabidopsis* development. RAP2 gene expression is both positively and negatively affected by the absence of AP2 activity during development. The observed differences in *RAP2.2*, *RAP2.3*, and *RAP2.4* gene expression levels in wild-type and *ap2-10* flowers and vegetative tissues are not apparently due to differences in ecotype because similar changes in gene expression levels were observed for all three RAP2 genes in stems when ecotype was controlled. The regulation of RAP2 gene expression by AP2 in stems clearly indicates that unlike other floral homeotic genes AP2 functions in both reproductive and vegetative development.

Example 5

This example shows that transgenic plants of the invention bear seed with altered fatty acid content and composition.

Antisense transgenic plants were prepared using *AP2*, *RAP2.8*, and *RAP2.1* (two independent plants) using methods described above. The fatty acid content and composition were determined using gas chromatography as described Broun and Somerville *Plant Physiol.* 113:933-942 (1997). The results are shown in Table VI (for *AP2*) and Table VII (for the *RAP2* genes). As can be seen there, the transgenic plants of the invention have increased fatty acid content as compared to wild-type plants. In addition, the profile of fatty acids is altered in the plants.

The results shown in Table VIII reveal that there is an approximately 7 mg of oil in a wild-type *Arabidopsis* seed. By contrast, there is approximately 9 mg in an *ap2-4* and *ap2-5* seed and approximately 14-15 mg in an *ap2-10* seed. In addition, the spectrum of fatty acids in wild-type and *ap2* mutant seeds are quantitatively indistinguishable. Thus, loss of AP2 activity increases total fatty acid content without detectable changes in fatty acid composition.

Table VI. Analysis of Arabidopsis seed fatty acid content and composition.

Plants	Seed Mass ($\mu\text{g}/\text{seed}$)	Total Fatty Acid Content ($\mu\text{g}/\text{seed}$)	Fatty acid methylesters (%) (area % by GLC)							
			16:0	18:0	18:1	18:2	18:3	20:1	22:1	24:0
<i>Mutant</i>										
<i>ap2-10</i>	27	3.94	9.4	4.4	2.5	70.8	0.8	6.6	0.1	1.9
<i>αAP2 transgenic plants</i>										
C24 15-522 (F1-1)	34	11.0	4.5	3.3	3.1	49.0	3.2	26.3	0.6	4.9
C34 15-542 (F1-2)	24	2.5	6.3	3.4	3.1	52.1	4.0	28.1	0.4	0.4
<i>Wild-type</i>										
C24	20	2.67	9.4	4.9	8.8	62.7	0.7	8.1	0.3	1.9

Table VII. Analysis of *Arabidopsis* RAP2 antisense transgenic seed fatty acid content and composition.

Plants	Seed Mass ($\mu\text{g}/\text{seed}$)	Total Fatty Acid Content ($\mu\text{g}/\text{seed}$)	16:0	18:0	18:1	18:2	18:3	20:1	22:1	24:0
<i>RAP2 antisense</i>										
<i>transgenic plants</i>										
RAP34-2.8C LE	24	9.7	5.1	3.0	3.6	53.6	3.2	29.3	0.3	0.2
RAP34-2.1A' COL	34	11.0	9.5	1.9	3.0	53.6	4.2	20.3	5.5	0.5
RAP34-2.1D' COL	26	8.1	4.6	2.6	2.6	50.3	3.2	33.2	0.3	1.0
<i>Wild-type</i>										
COL ^a	24	ND	6.0	4.0	14.0	27.0	18.0	22.0	2.0	ND

ND, not determined

^aPatak et al. (1994) Oil content and fatty acid composition of seeds of various ecotypes of *Arabidopsis thaliana*: a search for useful genetic variants. *Curr. Sci.* 67, 470-472.

Table VI Genetic control of total seed fatty acid content in *Arabidopsis* using AP2.

Genotype	Average seed mass (µg per seed)	Total seed fatty acid content ^a (µg per seed)	Percent increase in seed mass ^b	FA content ^b	Percent fatty acid methyl esters (area % by GC)							
					16:0	18:0	18:1	18:2	18:3	20:0	20:1	24:0
Ler ^c	24	7.3 (0.2)	*		6.0	4.1	22.7	26.7	17.4	2.2	19.3	1.6
ap2-4	33	9.1 (0.7)	+38%	+25%	6.6	3.7	16.5	26.9	23.7	2.8	19.6	2.8

Col-O ^d	22	6.8 (0.7)	*		8.1	4.0	13.6	33.3	20.9	2.1	16.6	1.4
ap2-5	29	9.2 (0.7)	+32%	+35%	6.9	4.1	14.8	30.7	20.8	2.2	17.0	2.2

C24 ^e	22	6.8 (0.1)	*		8.0	5.7	15.1	26.5	22.5	3.0	17.6	1.6
ap2-10 ^f	48	14.5 (0.9)	+109%	+113%	7.9	4.6	12.1	25.2	24.3	3.3	19.5	2.5
ap2-10 (+/-)	29	8.7 (0.7)	+32%	+28%	7.0	5.2	15.1	26.5	23.2	3.1	18.2	1.8

^aStandard deviation values are given in parentheses.^bValues for wild-type seed fatty acid content shown here and used for comparison were obtained using seed lots chosen according to the date of seed harvest to account for seasonal differences.^cControl seed ecotype Landsberg *erecta*^dControl seed ecotype Columbia^eControl seed ecotype C24^fTransgenic *ap2* mutant line in C24 background

Example 6

This example describes construction of promoter construct which are used to prepare expression cassettes useful in making transgenic plants of the invention. In particular, this example shows use of two preferred promoters, the promoter from the *AP2* gene and the promoter from the *Bell* gene.

Figure 5 shows a *AP2* promoter construct. pAP2 represents the 16.3 kb *AP2* promoter vector cassette that is used to generate chimeric genes for use in plant transformations described here. pAP2 is comprised of the 4.0 kb promoter region of the *Arabidopsis AP2* gene. The Ti plasmid vector used is pDE1000 vector (Plant Genetic Systems, Ghent, Belgium). The pDE1000 vector DNA was linearized with BamH1 and the *AP2* promoter region inserted as a 4.0 kb BamH1 DNA fragment from plasmid subclone pLE7.2. At the 3' end of the inserted *AP2* promoter region, designated *AP2*, lie three restriction sites (EcoR1, Sma1 and SnaB1) into which different gene coding regions can be inserted to generate chimeric *AP2* promoter/gene cassettes. NOS::NPTII represents the plant selectable marker gene NPTII under the direction of the nopaline synthase promoter which confers resistance to the antibiotic kanamycin to transformed plants cells carrying an integrated *AP2* promoter cassette. LB and RB represent the T-DNA left and right border sequences, respectively, that are required for transfer of T-DNA containing the *AP2* promoter cassette into the plant genome. PVS1 designates the bacterial DNA sequences that function as a bacterial origin of replication in both *E. coli* and *Agrobacterium tumefaciens*, thus allowing pAP2 plasmid replication and retention in both bacteria. Amp^R and Sm/Sp^R designate bacterial selectable marker genes that confer resistance to the antibiotics ampicillin and streptomycin/spectinomycin, respectively, and allows for selection of *Agrobacterium* strains that carry the pAP2 recombinant plasmid.

Figure 6 shows a *BEL1* promoter construct. pBEL1 represents the 16.8 kb *BEL1* promoter vector cassette that is used to generate chimeric genes for use in plant transformations described here. pBEL1 is comprised of the 4.5 kb promoter region of the *Arabidopsis BEL1* gene. The Ti plasmid vector used is pDE1000 vector (Plant Genetic Systems, Ghent, Belgium). The pDE1000 vector DNA was linearized will BamH1 and the *BEL1* promoter region inserted as a 4.5 kb BamH1-Bgl2 DNA fragment from plasmid subclone pλ1C9R (L. Reiser, unpublished). At the 3' end of the inserted *BEL1* promoter region, designated *BEL1*, lie three restriction sites (EcoR1, Sma1 and

SnaB1) into which different gene coding regions can be inserted to generate chimeric BEL promoter/gene cassettes. NOS::NPTII represents the plant selectable marker gene NPTII under the direction of the nopaline synthase promoter which confers resistance to the antibiotic kanamycin to transformed plants cells carrying an integrated BEL promoter cassette. LB and RB represent the T-DNA left and right border sequences, respectively, that are required for transfer of T-DNA containing the BEL promoter cassette into the plant genome. PVS1 designates the bacterial DNA sequences that function as a bacterial origin of replication in both *E. coli* and *Agrobacterium tumefaciens*, thus allowing pBEL plasmid replication and retention in both bacteria. Amp^R and Sm/Sp^R designate bacterial selectable marker genes that confer resistance to the antibiotics ampicillin and streptomycin/spectinomycin, respectively, and allows for selection of *Agrobacterium* strains that carry the pBEL recombinant plasmid.

Example 7

This example shows that transgenic plants of the invention have increased seed yield.

It is widely known that seed filling and the deposition of total seed contents is determined in part by the availability and supply of carbon- and nitrogen-containing compounds or assimilates to the developing seed. Thus, an increase in seed size and seed contents typically result from a decrease in total seed number in the presence of a fixed supply of photoassimilates. Since total seed number is dependent on many factors including both male and female fertility and since *ap2* mutations affect both ovule and stamen development, total seed number and total seed yield was measured in transgenic plants of the invention and in wild-type plants to determine whether the increase in seed size results at the expense of total seed number or seed yield.

To test this hypothesis directly, total seed yield for individual *ap2-10* (+/-) plants was determined. As shown in Table IX total seed yield per *ap2-10* (+/-) plant is increased by 35% when compared to wild-type C24, due in part to an increase in average seed mass (Table IX). In addition, increases in total seed yield in *ap2-10* transgenic plants may result from an increase in the total number of flowers produced. Table X shows that, on average, *ap2-10* (-/-) plants produce at least 80% more flowers than wild-type. Thus genetically manipulating AP2 activity in transgenic plants allows for

agriculturally desirable increases in total seed yield by increasing seed mass, seed contents, and number of flowers produced.

Table IX. Genetic control of Arabidopsis seed yield by AP2.

	Average seed mass (mg per 100 seed)	Total seed yield (g) ¹	Percent change in yield compared to wild-type
1. <i>ap2-10</i> (+/-)	2.9	2.61 (0.39)	+35%
2. C24	2.1	1.94 (0.27)	

¹n = 10. Standard deviation values are given in parentheses.

Table X. Genetic control of Arabidopsis flower number by AP2.

	Total # of inflorescences (per plant) ¹	Average number of flowers on primary inflorescence (per plant) ¹
1. <i>ap2-10</i> (-/-)	14.7 (3.9)	78.7 (6.0)
2. C24	11.0 (3.6)	42.9 (7.6)

¹n = 7. Standard deviation values are given in parentheses.

The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference for all purposes.

WHAT IS CLAIMED IS:

- 1 1. A method of modulating seed mass in a plant, the method
2 comprising:
3 providing a first plant comprising a recombinant expression cassette
4 containing an *ADC* nucleic acid linked to a plant promoter;
5 selfing the first plant or crossing the first plant with a second plant,
6 thereby producing a plurality of seeds; and
7 selecting seed with altered mass.
- 1 2. The method of claim 1, wherein expression of the *ADC* nucleic
2 acid inhibits expression of an endogenous *ADC* gene and the step of selecting includes
3 the step of selecting seed with increased mass.
- 1 3. The method of claim 2, wherein the seed have increased protein
2 content, carbohydrate content, or oil content.
- 1 4. The method of claim 2, wherein the *ADC* nucleic acid is linked to
2 the plant promoter in the antisense orientation.
- 1 5. The method of claim 2, wherein the *ADC* nucleic acid hybridizes
2 under stringent conditions to a nucleic acid having a sequence as set forth in Genbank
3 Accession Nos. U12546, AF003101, and AF003094.
- 1 6. The method of claim 2, wherein the *ADC* nucleic acid hybridizes
2 under stringent conditions to a nucleic acid having a sequence as set forth in SEQ ID
3 NO:1.
- 1 7. The method of claim 2, wherein the *ADC* nucleic acid hybridizes
2 under stringent conditions to a nucleic acid having a sequence as set forth in SEQ ID
3 NO:2.

1 8. The method of claim 2, wherein the *ADC* nucleic acid is selected
2 from a group consisting of Genbank accession numbers U12546, AF003094,
3 AF003095, AF003096, AF003097, AF003098, AF003099, AF003100, AF003101,
4 AF003102, AF003103, AF003104, and AF003105.

1 9. The method of claim 2, wherein the first and second plants are
2 the same species.

1 10. The method of claim 2, wherein the first and second plants are
2 members of the family Brassicaceae.

1 11. The method of claim 2, wherein the first and second plants are
2 members of the family Solanaceae.

1 12. The method of claim 2, wherein the plant promoter is a
2 constitutive promoter.

1 13. The method of claim 12, wherein the promoter is a CaMV 35S
2 promoter.

1 14. The method of claim 2, wherein the promoter is a tissue-specific
2 promoter.

1 15. The method of claim 14, wherein the promoter is ovule-specific.

1 16. A seed produced by the method of claim 2.

1 17. The method of claim 1, wherein expression of the *ADC* nucleic
2 acid enhances expression of an endogenous *ADC* gene and the step of selecting
3 includes the step of selecting seed with decreased mass.

1 18. The method of claim 17, wherein the *ADC* nucleic acid
2 hybridizes under stringent conditions to a nucleic acid having a sequence as set forth in
3 Genbank Accession Nos. U12546, AF003101, and AF003094.

1 19. The method of claim 17, wherein the *ADC* nucleic acid
2 hybridizes under stringent conditions to a nucleic acid having a sequence as set forth in
3 SEQ ID NO:1.

1 20. The method of claim 17, wherein the *ADC* nucleic acid
2 hybridizes under stringent conditions to a nucleic acid having a sequence as set forth in
3 SEQ ID NO:2.

1 21. The method of claim 17, wherein the *ADC* nucleic acid is
2 selected from a group consisting of Genbank accession numbers U12546, AF003094,
3 AF003095, AF003096, AF003097, AF003098, AF003099, AF003100, AF003101,
4 AF003102, AF003103, AF003104, and AF003105.

1 22. The method of claim 17, wherein the first and second plants are
2 the same species.

1 23. The method of claim 17, wherein the first and second plants are
2 members of the family Brassicaceae.

1 24. The method of claim 17, wherein the first and second plants are
2 members of the family Solanaceae.

1 25. The method of claim 17, wherein the plant promoter is a
2 constitutive promoter.

1 26. The method of claim 25, wherein the promoter is a CaMV 35S
2 promoter.

1 27. The method of claim 17, wherein the promoter is a tissue-specific
2 promoter.

1 28. The method of claim 27, wherein the promoter is ovule-specific.

1 29. A seed produced by the method of claim 17.

1 30. A seed comprising a recombinant expression cassette containing
2 an ADC nucleic acid.

1 31. The seed of claim 30, which is derived from a plant that is a
2 member of the family Brassicaceae.

1 32. The seed of claim 30, wherein the *ADC* nucleic acid hybridizes
2 under stringent conditions to a nucleic acid having a sequence as set forth in Genbank
3 Accession Nos. U12546, AF003101, and AF003094.

1 33. The seed of claim 30, wherein the *ADC* nucleic acid hybridizes
2 under stringent conditions to a nucleic acid having a sequence as set forth in SEQ ID
3 NO:1.

1 34. The seed of claim 30, wherein the *ADC* nucleic acid hybridizes
2 under stringent conditions to a nucleic acid having a sequence as set forth in SEQ ID
3 NO:2.

1 35. The seed of claim 30, wherein the *ADC* nucleic acid is selected
2 from a group consisting of Genbank accession numbers U12546, AF003094,
3 AF003095, AF003096, AF003097, AF003098, AF003099, AF003100, AF003101,
4 AF003102, AF003103, AF003104, and AF003105.

1 36. The seed of claim 30, wherein the *ADC* nucleic acid is linked to
2 a plant promoter in an antisense orientation and the seed mass is at least about 10%

3 greater than the average mass of seeds from the same plant variety which lack the
4 recombinant expression cassette.

1 37. The seed of claim 36, wherein the mass is at least about 20%
2 greater than the average mass of seeds from the same plant variety which lack the
3 recombinant expression cassette.

1 38. The seed of claim 36, wherein the mass is at least about 50%
2 greater than the average mass of seeds from the same plant variety which lack the
3 recombinant expression cassette.

1 39. The seed of claim 36, wherein the oil content is proportionally
2 increased.

1 40. The seed of claim 36, wherein the protein content is
2 proportionally increased.

1 41. The seed of claim 30, wherein the *ADC* nucleic acid is linked to
2 a plant promoter in the sense orientation and the seed mass is at least about 10% less
3 than the average mass of seeds of the same plant variety which lack the recombinant
4 expression cassette.

1 42. The seed of claim 41, which has a mass at least about 20% less
2 than the average mass of seeds of the same plant variety which lack the recombinant
3 expression cassette.

1 43. The seed of claim 41, which has a mass at least about 50% less
2 than the average mass of seeds of the same plant variety which lack the recombinant
3 expression cassette.

1 44. A transgenic plant comprising an expression cassette containing a
2 plant promoter operably linked to a heterologous *ADC* polynucleotide.

1 45. The transgenic plant of claim 44, wherein the *ADC* nucleic acid
2 hybridizes under stringent conditions to a nucleic acid having a sequence as set forth in
3 Genbank Accession Nos. U12546, AF003101, and AF003094.

1 46. The transgenic plant of claim 44, wherein the *ADC* nucleic acid
2 hybridizes under stringent conditions to a nucleic acid having a sequence as set forth in
3 SEQ ID NO:1.

1 47. The transgenic plant of claim 44, wherein the *ADC* nucleic acid
2 hybridizes under stringent conditions to a nucleic acid having a sequence as set forth in
3 SEQ ID NO:2.

1 48. The transgenic plant of claim 44, wherein the *ADC*
2 polynucleotide is selected from a group consisting of Genbank accession numbers
3 U12546, AF003094, AF003095, AF003096, AF003097, AF003098, AF003099,
4 AF003100, AF003101, AF003102, AF003103, AF003104, and AF003105.

1 49. The transgenic plant of claim 44, wherein the heterologous *ADC*
2 polynucleotide encodes a *ADC* polypeptide.

1 50. The transgenic plant of claim 44, wherein the heterologous *ADC*
2 polynucleotide is linked to the promoter in an antisense orientation.

1 51. The transgenic plant of claim 44, which is a member of the genus
2 *Brassica*.

1 52. An isolated nucleic acid molecule comprising an expression
2 cassette containing a plant promoter operably linked to a heterologous *ADC*
3 polynucleotide.

1 53. The isolated nucleic acid molecule of claim 52, wherein the *ADC*
2 nucleic acid hybridizes under stringent conditions to a nucleic acid having a sequence
3 as set forth in Genbank Accession Nos. U12546, AF003101, and AF003094.

1 54. The isolated nucleic acid molecule of claim 52, wherein the *ADC*
2 nucleic acid hybridizes under stringent conditions to a nucleic acid having a sequence
3 as set forth in SEQ ID NO:1.

1 55. The isolated nucleic acid molecule of claim 52, wherein the *ADC*
2 nucleic acid hybridizes under stringent conditions to a nucleic acid having a sequence
3 as set forth in SEQ ID NO:2.



1 56. The isolated nucleic acid molecule of claim 52, wherein the *ADC*
2 polynucleotide is selected from a group consisting of Genbank accession numbers
3 U12546, AF003094, AF003095, AF003096, AF003097, AF003098, AF003099,
4 AF003100, AF003101, AF003102, AF003103, AF003104, and AF003105.

1 57. The isolated nucleic acid of claim 52, wherein the heterologous
2 *ADC* polynucleotide encodes a *ADC* polypeptide.

1 58. The isolated nucleic acid of claim 52, wherein the heterologous
2 *ADC* polynucleotide is linked to the promoter in an antisense orientation.

1 59. The isolated nucleic acid of claim 52, which is a member of the
2 genus *Brassica*.

Figure 1. The AP2 domain






Figure 3. Schematic map of 35S-AP2 sense construct




Figure 2. Schematic map of 35S-AP2 antisense construct




Figure 4




Figure 5. Schematic diagram of pAP2.





Figure 6. Schematic diagram of pBEL1.



Regulation of gene expression in a eukaryotic cell.

Figure 7

SEQUENCE LISTING

<110> Jofuku, K. Diane
Okamuro, Jack K.
The Regents of the University of California

<120> Methods for Improving Seeds

<130> 023070-067230PC

<140> WO PCT/US99/03429
<141> 1999-02-17

<150> US 09/026,039
<151> 1998-02-19

<160> 104

<170> PatentIn Ver. 2.0

<210> 1
<211> 1669
<212> DNA
<213> Brassica napus

<220>
<223> canola (rape) APETALA2 (AP2) domain containing
(ADC) gene sequence

<220>
<221> modified_base
<222> (5)
<223> n = unknown

<220>
<221> modified_base
<222> (7)
<223> n = unknown

<220>
<221> modified_base
<222> (15)
<223> n = unknown

<220>
<221> modified_base
<222> (17)
<223> n = unknown

<220>
<221> modified_base
<222> (20)
<223> n = unknown

<220>
<221> modified_base
<222> (70)
<223> n = unknown

<220>
 <221> modified_base
 <222> (435)
 <223> n = unknown

<220>
 <221> modified_base
 <222> (1051)
 <223> n = unknown

<400> 1
 cagcngngtt ttcnntnatn gttcgtggcg gccacagtgg taggagaaga cggaaattaa 60
 caaattcatn gtcaccctac aagaaagggg gaacataatt aaatttcgag tagttggagt 120
 aggagaagct caatagtaca agaaattaaa taagatactc ccatctacat catcttgctt 180
 tttctcatcc cataatagtt ttccagtaaa actgtgaacc ttgtgaattt aatttcctt 240
 ttatatataa agataccttg tgggtgtttat actcaggaga ccagaaacta ggaaacagtc 300
 tattacattt gtattgggaa ataatacaat ctcaaaattt gattcattca taaactttat 360
 actatatact ttttcgttga taaatttttt gctctcctt ctaataacga atggagtcct 420
 agcacatata tattncctaa tgtgattttc atttcatatg gatattttat tatatatgac 480
 tattttattcc atcccatctt attgcttttg atggttctct tgtcattaga gtcttctctc 540
 ttattgctct catatttctt tgcttttgtt tcctctttat tacaagagag atatgtggaa 600
 ccttaacgac tcacctgatc accacgaaga atccgacggt agatggaaac gggctggaga 660
 tgtaccaatc tctatgagat catcgaccac gtgtctgtcg tctgttctc cctgacctcg 720
 gatatttttt tccgaatcaa atcatggaac aggaagttcc aggaatatct gggctccgta 780
 tcactagaaa ccagtctctt gttcggtcga atcctagcgg gtctggctcg cgggaaacc 840
 tagagctgga gataacacag ccggtaaaaa agagccgacg tggctctcgc tcacggagct 900
 ctgagtatag aggagttact ttttatcgac gaaccggaag atgggagtca catatttggt 960
 aacttaattt tcttaaccgg acgatatacc gaatactatt attacctata tggtaaactct 1020
 atcaaataca tgtttcattt catttgagcc nataccgtat tgttgttttt aaaatatgtt 1080
 tggaatctta tgcagggact gcggaagca agtgactta ggtatgatca tgtaatgttg 1140
 ttcaaacaca gatcaaatat cctattgaaa ctaagttgtg ttgtgtctgt ccatttttat 1200
 atgatttctt cgaccaaata aaggttttat tatctcctta tattactttt tgttacatat 1260
 tcaggtggat ttgacacagc acatgccgct gctcggtagt ttttactcat ccaaatatga 1320
 tcaattagaa cgaatctaatt attccttatt ttgtaatttg ctgatataca aattaatttg 1380
 ggtgggtaac tgtttgggac agtgccctac atagagccgc agttaagttt agagggtgag 1440
 atgcagatat aaatttcaat attgaagact atgtggagga tttgaaacag gtaaaatatt 1500
 tattatttgt aggttcaagc aattgactta gattattact cgaacataaa acaaatatt 1560
 atttggttga gatgacgcag ttgacaaagg aagagttcat gcatgtcatt agaaggcaaa 1620
 gcactgggtt tccaagaggc agctctaagt atagaggtgt cactttgca 1669

<210> 2
 <211> 803
 <212> DNA
 <213> Glycine max

<220>
 <223> soybean APETALA2 (AP2) domain containing (ADC)
 gene sequence

<400> 2
 ggattgtggg aaacaagttt atctaggtaa agttgattaa taacaataat tgtatatgtg 60
 tttgtgagaa ctgtggcagt ttttttctt aatattgttt taagaggcta aaacggtttt 120
 tttttccttg ttttgtgttt tttgtcttg ctgtgatgcg gtagagacaa gagtgtgagt 180
 gtgtgttgtg tgtgggtgag gatttttttr tttttttgtg gtgactgact tgatggtttt 240
 tgtctgggta aaatttgcct aggtggattt gacacagcac atgcggctgc tcggtgagcc 300
 cttgccccct cctttagtat tataccaagc tfgtaaatatt actttttcca tgtcttgaac 360
 caaatatcaa atattattgt aaatcacatt tcgttgtggg ccggggaatt gtgagtctca 420
 aagaaaattg tgtattttcc gtctctcttt tcagtgttta tgatagagcg gctattaaat 480
 tccgaggagt ggaggctgac attaaactca atattggaga ctatgaagat gacttgaagc 540
 aggtgatcaa tttgtggatt atgttttttt tattcgatat aaatgcattt atcgatttta 600
 tcttatcttg aacagtcata cgtataggat gcaccttato tcctacagtt agtgtttttt 660

```

tttatcttga attattctca tgattttgtt aaatgcaatg ttaatagatg agcaatctta 720
ccaaggaaga gttcgtccac gtgcttcgcc gccaaagcac tggatttccg agaggaagct 780
ccaagtatag aggtgtcact tgc 803

```

<210> 3
 <211> 11721
 <212> DNA
 <213> Arabidopsis thaliana

<220>
 <223> Arabidopsis APETALA2 (AP2) complete genomic
 sequence

<400> 3
 gtaagcaata atatagtttg aacaatgagt tcaaaagcct ttggttgga taactcatta 60
 agaaaacaga tagtttaaaa tcaattgaat tatgatttca gacatagtat aagtttgggc 120
 aaagtatttt aatggaaata gaatacaaac ttgataacaa gttattgcca tttagaaaat 180
 gttataactc cttatatagt gagaacccta ttgcttgctt gtgttcaagg aattgctgtt 240
 tatggaactc cactttaaat aaacatataa ataacttatt tagtatcaaa aacttaatat 300
 cccatctact ttgaatttgc ccgatcatga aaacacctat ttattgtaca taaatataca 360
 tctttaacaa atatagatct atttgtattt gtatcttatt accttttatt catatagaaa 420
 aaggaaaaaa caaacaacaa gctttgttcc aagagtttaa agtatatcaa aattggcagt 480
 attgtggggt ttgaagtaa tctactagat tattcatttt ctgcaaaaa acacctttt 540
 atagttgctt accgaaccaa gccgtggact tcactttctt aaaacatgaa attacatatt 600
 aaaggaatct ttgattaat caaagaagat gtttgtatac atagataata aatgaaagt 660
 gaaatatttt tttttaaaat aaagaaaaaa aataaagaaa ctccaatacg aggaattgtt 720
 tcgaatataa tcttttttcc aaagaaagca agtggttaga gagagaataa tattatttta 780
 ttaaattttt acaaaaaaaa cagtgaacag agaagagaga gagagaaagg gcagtggaag 840
 taaatataaa ggaaaggata aaaatgaaag ctttcgtaga agcaatctat caaattttta 900
 ttttattttt ttctctctct ctctttagct cttttttttt tgttttcatt aaagttttta 960
 ttttattttt taccaaccaa aagcttttct ctttggtttc tcttatttag cttctaacct 1020
 tgaggagaat cataccagag gattgaagtt tgaaccttca aagatcaaaa tcaagaaacc 1080
 aaaaaaaaaa aaaaaaaatg tgggatctaa acgacgcacc acaccaaaca caaagagaag 1140
 aagaatctga agagttttgt tattcttcac caagtaaacg gggtggatct ttctctaatt 1200
 ctactctctc agctgttgtt atcgaagatg gatccgatga cgatgaactt aaccgggtca 1260
 gacccaataa cccacttgct acccatcagt tcttccctga gatggattct aacggcggtg 1320
 gtgttgcttc tggctttcct cgggctcact gggttggtgt taagttttgt cagtcggatc 1380
 tagccaccgg atcgtccgag ggtaaagcta ccaacgttgc cgctgccgta gtggagccgg 1440
 cacagccgtt gaaaaagagt cggcgtggac caagatcaag aagttctcag tatagaggtg 1500
 ttacgtttta ccggcgatcc ggaagatggg aatctcatat ttggtaataa tctcatattt 1560
 ttaatttcgt taatcgatcg tacttttagat tataaattta agtttttttt tgtttgttct 1620
 tctgaatttc agggactgtg ggaaacaagt ttacttaggt aattttattt tcctcatgtt 1680
 tttttttgta ttttggtgtt gaaaaatgtc atcataattt taatttatta taanctctga 1740
 ataggtggat ttgacactgc tcatgcagca gctcggatatt tttctctctt tgactctctc 1800
 tatattgagt tgttatttat ttattttttt aaaaataccg gaagaaattt ataaaaatta 1860
 attttaattt tgttttattt aatagagcat atgatagagc tgctattaaa ttccgtggag 1920
 tagaagcggg tatcaatttc aacatcgaag attatgatga tgacttgaaa caggtaaata 1980
 taaattataa actatatttg tttttattaa cgatttttaa aggtttggga gattaatatt 2040
 gaaattgaat tttatagatg actaatttaa ccaaggaaga gttcgtacac gtacttcgcc 2100
 gacaaagcac aggtctccct cgaggaagtt cgaagtatag aggtgtcact ttgcataagt 2160
 gtggtcgttg ggaagctcga atgggtcaat tcttaggcaa aaagtataat ttctctcatt 2220
 ttatattcac tcgaaaactt catttttagt ttgttatttt aactttgagt ttttgtttct 2280
 tgaatcttat aaaaataggta tgtttatttg gggttggtcg acaccgaggt cgaagctgct 2340
 aggtaaatgt ctttttggtt gattctacaa cacacattgt tgtataatgt gtttttctcg 2400
 ttactaatgt attttcatta ttttatatat aatcacagag cttacgataa agctgcaatc 2460
 aatgtaacg gcaaaagcgc cgtgaccaac ttgatccga gtatttacga tgaggaactc 2520
 aatgccggta aattgtctca tttaatcgag taattttata tatttttttg tccttagttt 2580
 catcactttg gtgttcgaac ttggttttaa gattttgaat ttggtgtata tagagtcatc 2640
 agggaatcct actactccac aagatcacia cctcgatttg agcttgggaa attcggctaa 2700
 ttcgaagcat aaaagtcaag atatgcggct cagggtaggg ttaaatctta tattattaac 2760
 aataatttat atcttaatat attgtttata tgtttataaa catgttttct tttgttttgc 2820

tttcagatga	accaacaaca	acaagattct	ctccactcta	atgaagttct	tggattaggt	2880
caaaccggaa	tgccttaacca	tactcccaat	tcaaaaccacc	aagtgaagtaa	ataaccacaa	2940
atgcaaatatc	cataatttca	tttgaatata	ttttatctaa	agaattgcat	ttttttttgg	3000
taaattagtt	tccgggcagc	agcaacattg	gtagcggagg	cggattctca	ctgtttccgg	3060
cggctgagaa	ccaccggttt	gatggtcggg	cctcgacgaa	ccaagtgttg	acaaatgctg	3120
cagcatcatc	aggattctct	cctcatcatc	acaatcagat	ttttaattct	acttctactc	3180
ctcatcaaaa	ttggctgcag	acaaatggct	tccaacctcc	tctcatgaga	ccttcttgaa	3240
tcttttata	ttttaagggt	tattattata	taagaaaaac	aaaaatgaac	ccttgaaatc	3300
cccacatgtt	cttggtcatt	tcattaatca	tcggccttata	ttttgcttat	tttcccctaa	3360
atcctcttgt	taacttaggc	gaacaaaaaa	aattaatgga	aatctttttc	cctccatcgg	3420
ttacaaaaat	aatattatat	ataattgttg	gatatatggg	aaattggata	agtttgtgat	3480
ttgagatgtt	ctgactaaaa	aggttgagaa	gagatttgtc	aatgagttgt	cctttgttgt	3540
ttctcctcaa	tacatttatt	aagattttta	aacattactt	ctttatatgt	tacctgcacc	3600
tatctacata	tatgttcagt	cctaacttgt	tttgttattc	ctctttcata	tctattcaat	3660
taatgttttc	ctagccttag	ttcattttac	atttttcttg	aaaatctctc	atgaaaaaaa	3720
cacattcatg	tgtgaaatat	atttcaacac	ccattatagt	ttcgtttaatt	cagatatata	3780
atttttattt	attacatata	ataaaaaattg	acaggtgggt	atacacatgg	tttccttgtg	3840
ctaacattgg	tttgaatagc	ataaacggaa	tcctaaaata	ctttaagttc	acttcgtaaa	3900
taaaatctga	taaactgaac	aaattaggtt	ctctaaagtt	gagatgggta	aatgttcagc	3960
taactcatgg	agttggaatt	gtgattctcg	cttggtcaca	attgcttttg	gtattcacaa	4020
ggatacta	tgtatgcttt	gtttgtggtt	gtgctttgtt	tgccattggt	actcctttat	4080
ggagatctat	gttctatctt	gtncctctgt	tgttgggaaa	tctttatttg	cagggaaatga	4140
aagcttcttt	ggtgtttatg	gatttgcgtg	tgcactctgt	ttggacatgc	tatcttctaa	4200
aacccttgaa	gttcctctgc	gagacatttc	tatcgcttat	cgttctagca	cctacacggt	4260
ggtagcagaa	gatattgtag	cgaaaactgc	cttatcattg	aataacgccc	tcttgtttggc	4320
gtggaacgaa	gactctttgg	ttttgcaatt	gttcatggac	tcccagttca	gtgtaaaatc	4380
tttcaagtca	ctcgaagtca	ccgttgagct	agggagaatt	ctgcttgttt	taagcttctt	4440
tcaccttcgg	tttatcccca	tgttttttgc	attattctct	tatactgcgt	ttgtgtttgc	4500
agttctttta	gttttttagtg	atgtgtttta	ctcttgtaat	ggagactaaa	ttatttctat	4560
atctataaaa	ttagtttgat	aaaaaaaaaa	aaactgaaca	aattttattt	cgttcagcaa	4620
cgcttataaa	tgccaaatag	aaaagttcag	ttatttttaa	gcttttttgc	atatatatat	4680
tatctaacct	tttattgaat	tagagaattt	tagtcaacca	ttaaaaatta	cgaattttcc	4740
tagtttatac	tatttagaag	ttgcagtatc	aatcatcacg	ggaaaaaaat	aaaatgatat	4800
ttttggtttt	gtccattttg	tagatatttt	accgacagaa	gaaaaaaaca	gtgatatttt	4860
atttttttac	agtattaatg	gtgagacgag	agagagaaac	agaagtagat	tggttcttat	4920
gtttccctaa	tggagtaaga	aataatactc	acagttattt	tgcgatctca	acaaaagtta	4980
aaatgaatat	caaccgtaag	atctctttct	tctgttttca	cagactatga	aatataaaaa	5040
atcataatgc	ctacaagcta	tgcgaccggt	agttaaaaaa	aaatatgaat	aatattccaa	5100
agagaaaaaa	atcttggaag	tagaaaatta	ggtataaaga	gaaaagaggc	aaataaaaaat	5160
tggggattat	tagggatgag	tggtcatttt	caagtaatgt	gtcctttgaa	tcagtttctc	5220
tctctatctc	tcacagaaac	ccaaaagaag	tcagacattg	ttataatggt	gagagagact	5280
tctcggtctc	acttcttttc	tctcttttta	attctctttt	taattcaaaa	gtttaaagat	5340
tttattaaat	gtttcatact	cttacactta	ttcatgaatc	ctttcctaga	tttcactttt	5400
actctctgta	taatttggtc	ctccttaaac	tcttggttca	tctatttggt	tacgttacat	5460
cctaatagtc	tttatcatat	ataccttttg	gtcctttttac	taatactagg	taaaaacttc	5520
tagttgtata	atgcttaaaa	ttacaatggg	gttatgatga	tattgttatt	ttataatcgc	5580
attagaattg	caaacaaaaa	ttgtttatgc	taggataaac	attttaagtg	agaagactat	5640
cccttttttt	atatctatat	ttagattcgg	aattttctta	ctaacgaaaa	atatatagat	5700
ggagaactat	gacaatacac	ttttgcctta	caatgacatt	atggttattt	acctaattga	5760
atgatacaaa	atgagatgga	gttgtatgaa	tttatagcaa	ctgtctttct	gcttcttttt	5820
ttttttttta	atagagtgga	gacttgaatt	cnnwrtnnag	natgnnnncy	gaattcaagt	5880
ctccactcta	ttaaaaaaaa	aaaaaagaag	cagaaagaca	gttgctataa	attcatacaa	5940
ctccatctca	ttttgtatca	ttcaattagg	taaataacca	taatgtcatt	gtaaggcaaa	6000
agtgtattgt	catagttctc	catctatata	tttttcgtta	gtaagaaaat	tccgaatcta	6060
aatatagata	taaaaaaagg	gatagtcttc	tcacttaaaa	tgtttatcct	agcataaaca	6120
atttttgttt	gcaattctaa	tgcgattata	aataaacaat	atcatcataa	ccccattgta	6180
attttaagca	ttatacaact	agaagttttt	acctagtatt	agtaaaagga	cccaaaggta	6240
tatatgataa	agactattag	gatgtaacgt	aaccaaatag	atgaaccaag	agtttaagga	6300
ggaacaaatt	atacagagag	taaaagtga	atctaggaaa	ggattcatga	ataagtgtaa	6360
gagtatgaaa	catttaataa	aatcttttaa	cttttgatt	aaaaagagaa	ttaaaagagg	6420
agaaaggaag	tgaagccgag	aagtctctct	caccattata	acaatgtctg	acttcttttg	6480

ggtttctgtg	agagatagag	agagaaactg	attcaaagga	cacattactt	gaaaatgacc	6540
actcatccct	aataatcccc	aattttttatt	tgcctctttt	ctctttatac	ctaattttct	6600
atttccaaga	tttttttctc	tttggaatat	tattcatatt	tttttttaac	taacgggtcg	6660
atagcttgta	ggcattatga	ttttttatat	ttcatagtct	gtgaaaacag	aagaaagaga	6720
tcttacgggt	gatattcatt	ttaacttttg	ttgagatcgc	aaaataactg	tgagtattat	6780
ttcttactcc	attagggaaa	cataagaacc	aatctacttc	tgtttctctc	tctcgtctca	6840
ccattaatac	tgtaaaaaaa	taaaatatca	ctgttttttt	cttctgtcgg	taaaatatct	6900
acaaaatgga	caaaacccaa	aatatcattt	tatttttttc	ccgtgatgat	tgatactgca	6960
acttctaaat	agtataaact	aggaaaattc	gtaattttta	atgggtgact	aaaattctct	7020
aattcaataa	aaggttagat	aatatatata	tgcaaaaaag	ctttaaaata	actgaacttt	7080
tctatttggc	atttataagc	gttgctgaac	gaaataaaat	ttgttcagtt	tttttttttt	7140
tatcaaacta	attttataga	tatagaaata	atttagtctc	cattacaaga	gtaaaacaca	7200
tcactaaaaa	ctgaaagaac	tgcaaacaca	aacgcagtat	aagagaataa	tgcaaaaaac	7260
atggggataa	accgaagggt	aaagaagctt	aaaacaagca	gaatttctcc	tagctcaacg	7320
gtgacttcga	gtgacttgaa	agattttaca	ctgaactggg	agtccatgaa	caattgcaaa	7380
accaaagagt	cttcgttcca	cgccaacaag	agggcggtat	tcaatgataa	ggcagttttc	7440
gctacaatat	cttctgtctac	caacgtgtag	gtgctagaac	gataagcgat	agaaatgtct	7500
cgcaagggaa	cttcaagggt	tttagaagat	agcatgtcca	aacaagatgc	aacagcaaat	7560
ccataaacac	caaagaagct	ttcattccct	gcaaataaag	atttcccaac	aacaggggna	7620
caagatagaa	catagatctc	cataaaggag	taacaatggc	aaacaaagca	caaccacaaa	7680
caaagcatac	aattagtatc	cttgtgaata	ccaaaagcaa	ttgtgaacaa	gcgagaatca	7740
caattccaac	tccatgagtt	agctgaacat	ttaccatctc	caactttaga	gaacctaat	7800
tggttcagttt	atcagatttt	atttacgaag	tgaacttaaa	gtatttttagg	attcgggttta	7860
tgctatttcaa	accaatgtta	gcacaaggaa	accatgtgta	taccacctg	tcaattttta	7920
ttatatgtaa	taaaataaaa	ttatatatct	gaattaacga	aactataatg	ggtgttgaaa	7980
tatatattcac	acatgaatgt	gtttttttca	tgagagattt	tcaagaaaaa	tgtaaaatga	8040
actaaggcta	ggaaaacatt	aattgaatag	atatgaaaga	ggaataacaa	aacaagttag	8100
gactgaacat	atatgtagat	aggtgcaggt	aacatataaa	gaagtaatgt	tttaaaatct	8160
taataaatgt	attgaggaga	aacaacaaaa	gacaactcat	tgacaaatct	cttctcaacc	8220
tttttagtca	gaacatctca	aatcacaaac	ttatccaatt	tcccatatat	ccaacaatta	8280
tatataatat	tattttttgta	accgatggag	ggaaaaagat	ttccattaat	tttttttggt	8340
cgcttaagtt	aacaagagga	tttaggggaa	aataagcaaa	atataagccg	atgattaatg	8400
aaatgaccaa	gaacatgtgg	ggatttcaaa	ggttcatttt	tgtttttctt	atataataat	8460
aaaccttaaa	aataataaag	attcaagaag	gtctcatgag	aggagggttg	aagccatttg	8520
tctgcagcca	attttgatga	ggagtagaag	tagaattaaa	aatctgattg	tgatgatgag	8580
gagagaatcc	tgatgatgct	gcagcatttg	tcaacacttg	gttcgtcgag	gcccgcacat	8640
caaaccgggtg	gttctcagcc	gccggaaaca	gtgagaatcc	gcctccgcta	ccaatgttgc	8700
tgctgcccgg	aaactaattt	acaaaaaaa	aatgcaattc	tttagataaa	atatattcaa	8760
atgaaattat	ggtatttgca	tttgtgggta	tttactcact	tgggtggttg	aattgggagt	8820
atgggttaagc	attccgggtt	gacctaatcc	aagaacttca	ttagagtggg	gagaatcttg	8880
ttgttggttg	ttcatctgaa	agcaaaaaca	aagaaaacat	gtttataaac	atataaaca	8940
tatattaaga	tataaattat	tgtaataaat	ataagattaa	accctaccct	gagccgcata	9000
tcttgacttt	tatgcttcga	attagccgaa	tttcccaagc	tcaaatcgag	gttgtgatct	9060
tgtggagtag	taggattccc	tgatgactct	atatacacca	aattcaaaat	ctttaaacca	9120
agttcgaaca	ccaaagtgat	gaaactaagg	acaaaaaaat	atataaaatt	actcgattaa	9180
atgagacaat	ttaccggcat	tgagttcctc	atcgtaaata	ctcgatcaa	agttggtcac	9240
ggcgtctttg	ccgttacatt	tgattgcagc	tttatcgtaa	gctctgtgat	tatatataaa	9300
ataatgaaaa	tcaattagta	acgagaaaaa	cacattatac	aacaatgtgt	gttgtagaat	9360
caaacaaaaa	gacatttacc	tagcagcttc	gacctcggtg	tcgaacaaac	ccaaataaac	9420
atacctattt	tataagattc	aagaaacaaa	aactcaaagt	taaaataaca	aactaaaaat	9480
gaagttttcg	agtgaatata	aaatgagaga	aattataact	tttgcctaag	aattgacca	9540
ttcgagcttc	ccaacgacca	cacttatgca	aagtgcacc	tctatacttc	gaacttcctc	9600
gagggaaagc	tgtgctttgt	cggcgaagta	cgtgtacgaa	ctcttccttg	gttaaatag	9660
tcatctataa	aattcaattt	caatattaat	ctcccaaac	tttaaaaatc	gttaataaaa	9720
accaatatag	tttataattt	atatttacct	gtttcaagtc	atcatcataa	tcttcgatgt	9780
tgaaattgat	atccgcttct	actccacgga	atttaatagc	agctctatca	tatgctctat	9840
taaaataaac	aaaatttaaa	ttaattttta	taaatttctt	ccggtatttt	taaaaaaata	9900
aataaataac	aactcaatat	agagagagtc	aaagagagaa	aaataccgag	ctgctgcag	9960
agcagtgta	aatccaccta	ttcagagntt	ataataaatt	aaaattatga	tgacattttt	10020
caacacccaa	atacaaaaaa	aaacatgagg	aaaataaaat	tacctaagta	aacttgtttc	10080
ccacagtccc	tgaaattcag	aagaacaaac	aaaaaaaaac	ttaaatttat	aacttaaggt	10140

```

acgatcgatt aacgaaatta aaaatatgag attattacca aatatgagat tcccatcttc 10200
cggtagcgcc gtaaaacgta acacctctat actgagaact tcttgatctt ggtccacgcc 10260
gactcttttt caacggctgt gccggctcca ctacggcagc ggcaacgttg gtagctttac 10320
ccgcggaagc tccggtgggt agatccgact gacaaaactt aacaccaaac cagtgaagccc 10380
gaggaaagcc agaagcaaca ccaccgccgt tagaatccat ctcagggaag aactgatggg 10440
tgacaagtgg gttattgggt ctgaccgggt taagttcatc gtcacggat ccatcttcga 10500
taacaacagc tgaagagcta gaattagaga aagatccaac ccgtttactt ggtgaagaat 10560
aacaaaactc ttcagattct tcttctcttt gtgtttggtg tgggtgcgtcg tttagatccc 10620
acattttttt tgtttttttt tggtttcttg attttgatct ttgaagggtc aaacttcaat 10680
cctctggtat gattctcctc aagggttagaa gctaaataag agaaaccaa gagaaaagct 10740
tttggttggg agaaaataaa ataaaaactt taatgaaaac aaaaaaaaa gagctaaaga 10800
gagagagaga agaaaataaa ataaaaattt gatagattgc ttctacgaaa gctttcattt 10860
ttatcttttc ctttatattt acttccactg ccttttctct ctctctcttc tcttgctact 10920
gttttttttg taaaaattta ataaaaataat attattctct ctctaaccac ttgctttctt 10980
tggaaaaaag attatattcg aaacaattcc tcgtattgga gtttctttat tttttttctt 11040
tattttaaaa aaaaatattt ccactttcat ttattatcta tgtatacaaa catcttcttt 11100
gattaatcaa aagattcctt taatatgtaa tttcatgttt taagaaagtg aagtcacagg 11160
cttggttcgg taagcaacta taaaagggtg ttttttgcaa gaaaatgaat aatctagtag 11220
attacttcaa aacccacaaa tactgccaat tttgatatac tttaaactct tggaaacaaag 11280
ctttttgttt gttttttcct ttttctatat gaataaaagg taataagata caaatacaaa 11340
tagatctata tttgttaaag atgtatatatt atgtacaata aatagggtgt ttcgatgcg 11400
ggcaaattca aagtagatgg gatattaagt ttttgatact aaatagatta tttatatgtt 11460
tatttaaagt ggagttccat aaacagcaat tccttgaaca caagcaagca atagggttct 11520
cactatataa ggagttataa cattttctaa atggcaataa cttgttatca agtttgatt 11580
ctatttccat taaaatactt tgcccaaact tatactatgt ctgaaatcat aattcaattg 11640
attttaaact atctgttttc ttaatgagtt attccaacca aaggcttttg aactcattgt 11700
tcaaactata ttattgctta c 11721

```

<210> 4

<211> 67

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> DOMAIN

<222> (1)..(67)

<223> AP2-R1 AP2-like subclass AP2 domain repeat, amino
acid positions 129-195

<400> 4

```

Ser Ser Gln Tyr Arg Gly Val Thr Phe Tyr Arg Arg Thr Gly Arg Trp
  1             5             10             15

```

```

Glu Ser His Ile Trp Asp Cys Gly Lys Gln Val Tyr Leu Gly Gly Phe
      20             25             30

```

```

Thr Asp Ala His Ala Ala Ala Arg Ala Tyr Asp Arg Ala Ala Ile Lys
    35             40             45

```

```

Phe Arg Gly Val Glu Ala Asp Ile Asn Phe Asn Ile Asp Asp Tyr Asp
    50             55             60

```

```

Asp Asp Leu
    65

```

<210> 5

<211> 68

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> DOMAIN

<222> (1)..(68)

<223> AP2-R2 AP2-like subclass AP2 domain repeat, amino acids 221-288

<400> 5

Ser	Ser	Lys	Tyr	Arg	Gly	Val	Thr	Leu	His	Lys	Cys	Gly	Arg	Trp	Glu
1					5				10					15	

Ala	Arg	Met	Gly	Gln	Phe	Leu	Gly	Lys	Lys	Tyr	Val	Tyr	Leu	Gly	Leu
			20					25					30		

Phe	Asp	Thr	Glu	Val	Glu	Ala	Ala	Arg	Ala	Tyr	Asp	Lys	Ala	Ala	Ile
		35						40				45			

Lys	Cys	Asn	Gly	Lys	Asp	Ala	Val	Thr	Asn	Phe	Asp	Pro	Ser	Ile	Tyr
	50					55					60				

Asp	Glu	Glu	Leu
65			

<210> 6

<211> 18

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> HELIX

<222> (1)..(18)

<223> putative AP2-R1 amphipathic alpha-helix, amino acids 160-177

<400> 6

Phe	Asp	Thr	Ala	His	Ala	Ala	Ala	Arg	Ala	Tyr	Asp	Arg	Ala	Ala	Ile
1					5				10					15	

Lys Phe

<210> 7

<211> 18

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> HELIX

<222> (1)..(18)

<223> putative AP2-R2 amphipathic alpha-helix, amino acids 253-270

<400> 7

Phe	Asp	Thr	Glu	Val	Glu	Ala	Ala	Arg	Ala	Tyr	Asp	Lys	Ala	Ala	Ile
1					5				10					15	

Lys Cys

<210> 8
 <211> 4
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:AP2-like
 subclass AP2 domain conserved RAYD element

<400> 8
 Arg Ala Tyr Asp
 1

<210> 9
 <211> 77
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(77)
 <223> ANT-R1 AP2-like subclass AP2 domain repeat

<400> 9
 Thr Ser Gln Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr
 1 5 10 15
 Glu Ala His Leu Trp Asp Asn Ser Phe Lys Lys Glu Gly His Ser Arg
 20 25 30
 Lys Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Met Glu Glu Lys Ala
 35 40 45
 Ala Arg Ala Tyr Asp Leu Ala Ala Leu Lys Tyr Trp Gly Pro Ser Thr
 50 55 60
 His Thr Asn Phe Ser Ala Glu Asn Tyr Gln Lys Glu Ile
 65 70 75

<210> 10
 <211> 69
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(69)
 <223> ANT-R2 AP2-like subclass AP2 domain repeat

<400> 10
 Ala Ser Ile Tyr Arg Gly Val Thr Arg His His Gln His Gly Arg Trp
 1 5 10 15
 Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys Asp Leu Tyr Leu Gly
 20 25 30
 Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala Tyr Asp Val Ala Ala
 35 40 45

Ile Lys Phe Arg Gly Thr Asn Ala Val Thr Asn Phe Asp Ile Thr Arg
 50 55 60

Tyr Asp Val Asp Arg
 65

<210> 11
 <211> 67
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(67)
 <223> RAP2.7-R1 AP2-like subclass AP2 domain repeat

<400> 11
 Ser Ser Gln Tyr Arg Gly Val Thr Phe Tyr Arg Arg Thr Gly Arg Trp
 1 5 10 15

Glu Ser His Ile Trp Asp Cys Gly Lys Gln Val Tyr Leu Gly Gly Phe
 20 25 30

Asp Thr Ala His Ala Ala Ala Arg Ala Tyr Asp Arg Ala Ala Ile Lys
 35 40 45

Phe Arg Gly Val Asp Ala Asp Ile Asn Phe Thr Leu Gly Asp Tyr Glu
 50 55 60

Glu Asp Met
 65

<210> 12
 <211> 53
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(53)
 <223> RAP2.7-R2 AP2-like subclass AP2 domain repeat

<400> 12
 Ser Ser Lys Tyr Arg Gly Val Thr Leu His Lys Cys Gly Arg Trp Glu
 1 5 10 15

Ala Arg Met Gly Gln Phe Leu Gly Lys Lys Ala Tyr Asp Lys Ala Ala
 20 25 30

Ile Asn Thr Asn Gly Arg Glu Ala Val Thr Asn Phe Glu Met Ser Ser
 35 40 45

Tyr Gln Asn Glu Ile
 50

<210> 13
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:AP2-like
subclass AP2 domain conserved YRG element YRG
motif consensus sequence

<400> 13
Tyr Arg Gly Val Thr
1 5

<210> 14
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:AP2-like
subclass AP2 domain conserved YRG element
WEAR/WESH motif consensus sequence

<220>
<221> MOD_RES
<222> (5)
<223> Xaa = Ala or Ser

<220>
<221> MOD_RES
<222> (6)
<223> Xaa = Arg or His

<400> 14
Gly Arg Trp Glu Xaa Xaa
1 5

<210> 15
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:AP2-like
subclass AP2 domain RAYD element consensus
sequence

<400> 15
Val Tyr Leu Gly
1

<210> 16
<211> 4
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:AP2-like
subclass AP2 domain conserved RAYD element
consensus sequence

<400> 16

Ala Ala Ile Lys
1

<210> 17

<211> 69

<212> PRT

<213> Nicotiana tabacum

<220>

<221> DOMAIN

<222> (1)..(69)

<223> EREBP-1 EREBP-like subclass AP2 domain

<400> 17

Gly Arg His Tyr Arg Gly Val Arg Arg Arg Pro Trp Gly Lys Phe Ala
1 5 10 15

Ala Glu Ile Arg Asp Pro Ala Lys Asn Gly Ala Arg Val Trp Leu Gly
20 25 30

Thr Tyr Glu Thr Asp Glu Glu Ala Ala Ile Ala Tyr Asp Lys Ala Ala
35 40 45

Tyr Arg Met Arg Gly Ser Lys Ala His Leu Asn Phe Pro Leu Glu Val
50 55 60

Ala Asn Phe Lys Gln
65

<210> 18

<211> 69

<212> PRT

<213> Nicotiana tabacum

<220>

<221> DOMAIN

<222> (1)..(69)

<223> EREBP-2 EREBP-like subclass AP2 domain

<400> 18

Gly Arg His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly Lys Phe Ala
1 5 10 15

Ala Glu Ile Arg Asp Pro Ala Lys Asn Gly Ala Arg Val Trp Leu Gly
20 25 30

Thr Tyr Glu Thr Ala Glu Glu Ala Ala Leu Ala Tyr Asp Lys Ala Ala
35 40 45

Tyr Arg Met Arg Gly Ser Lys Ala Leu Leu Asn Phe Pro His Arg Ile
50 55 60

Gly Leu Asn Glu Pro
65

<210> 19
<211> 68
<212> PRT
<213> Nicotiana tabacum

<220>
<221> DOMAIN
<222> (1)..(68)
<223> EREBP-3 EREBP-like subclass AP2 domain

<400> 19
Glu Val His Tyr Arg Gly Val Arg Lys Arg Pro Trp Gly Arg Tyr Ala
1 5 10 15

Ala Glu Ile Arg Asp Pro Gly Lys Lys Ser Arg Val Trp Leu Gly Thr
20 25 30

Phe Asp Thr Ala Glu Glu Ala Ala Lys Ala Tyr Asp Thr Ala Ala Arg
35 40 45

Glu Phe Arg Gly Pro Lys Ala Lys Thr Asn Phe Pro Ser Pro Thr Glu
50 55 60

Asn Gln Ser Pro
65

<210> 20
<211> 69
<212> PRT
<213> Nicotiana tabacum

<220>
<221> DOMAIN
<222> (1)..(69)
<223> EREBP-4 EREBP-like subclass AP2 domain

<400> 20
Lys Lys His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly Lys Phe Ala
1 5 10 15

Ala Glu Ile Arg Asp Pro Asn Arg Lys Gly Thr Arg Val Trp Leu Gly
20 25 30

Thr Phe Asp Thr Ala Ile Glu Ala Ala Lys Ala Tyr Asp Arg Ala Ala
35 40 45

Phe Lys Leu Arg Gly Ser Lys Ala Ile Val Asn Phe Pro His Arg Ile
50 55 60

Gly Leu Asn Glu Pro
65

<210> 21
 <211> 68
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(68)
 <223> RAP2.2 EREBP-like subclass AP2 domain

<400> 21
 Lys Asn Gln Tyr Arg Gly Ile Arg Gln Arg Pro Trp Gly Lys Trp Ala
 1 5 10 15
 Ala Glu Ile Arg Asp Pro Arg Lys Gly Ser Arg Glu Trp Leu Gly Thr
 20 25 30
 Phe Asp Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Ala Ala Ala Arg
 35 40 45
 Arg Ile Arg Gly Thr Lys Ala Lys Val Asn Phe Pro Glu Glu Lys Asn
 50 55 60
 Pro Ser Val Val
 65

<210> 22
 <211> 68
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(68)
 <223> RAP2.3 EREBP-like subclass AP2 domain

<400> 22
 Lys Asn Val Tyr Arg Gly Ile Arg Lys Arg Pro Trp Gly Lys Trp Ala
 1 5 10 15
 Ala Glu Ile Arg Asp Pro Arg Lys Gly Val Arg Val Trp Leu Gly Thr
 20 25 30
 Phe Asn Thr Ala Glu Glu Ala Ala Met Ala Tyr Asp Val Ala Ala Lys
 35 40 45
 Gln Ile Arg Gly Asp Lys Ala Lys Leu Asn Phe Pro Asp Leu His His
 50 55 60
 Pro Pro Pro Pro
 65

<210> 23
 <211> 68
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(68)
 <223> RAP2.5 EREBP-like subclass AP2 domain

 <400> 23
 Glu Ile Arg Tyr Arg Gly Val Arg Lys Arg Pro Trp Gly Arg Tyr Ala
 1 5 10 15
 Ala Glu Ile Arg Asp Pro Gly Lys Lys Thr Arg Val Trp Leu Gly Thr
 20 25 30
 Phe Asp Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Thr Ala Ala Arg
 35 40 45
 Asp Phe Arg Gly Ala Lys Ala Lys Thr Asn Phe Pro Thr Phe Leu Glu
 50 55 60
 Leu Ser Asp Gln
 65

<210> 24
 <211> 68
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(68)
 <223> RAP2.6 EREBP-like subclass AP2 domain

 <400> 24
 Pro Lys Lys Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly Lys Trp Ala
 1 5 10 15
 Ala Glu Ile Arg Asp Pro His Lys Ala Thr Arg Val Trp Leu Gly Thr
 20 25 30
 Phe Glu Thr Ala Glu Ala Ala Ala Arg Ala Tyr Asp Ala Ala Ala Leu
 35 40 45
 Arg Phe Arg Gly Ser Lys Ala Lys Leu Asn Phe Pro Glu Asn Val Gly
 50 55 60
 Thr Gln Thr Ile
 65

<210> 25
 <211> 68
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> DOMAIN
 <222> (1)..(68)
 <223> RAP2.12 EREBP-like subclass AP2 domain

<400> 25

Lys Asn Gln Tyr Arg Gly Ile Arg Gln Arg Pro Trp Gly Lys Trp Ala
 1 5 10 15

Ala Glu Ile Arg Asp Pro Arg Glu Gly Ala Arg Ile Trp Leu Gly Thr
 20 25 30

Phe Lys Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Ala Ala Ala Arg
 35 40 45

Arg Ile Arg Gly Ser Lys Ala Lys Val Asn Phe Pro Glu Glu Asn Met
 50 55 60

Lys Ala Asn Ser
 65

<210> 26

<211> 68

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> DOMAIN

<222> (1)..(68)

<223> TINY EREBP-like subclass AP2 domain

<400> 26

His Pro Val Tyr Arg Gly Val Arg Lys Arg Asn Trp Gly Lys Trp Val
 1 5 10 15

Ser Glu Ile Arg Glu Pro Arg Lys Lys Ser Arg Ile Trp Leu Gly Thr
 20 25 30

Phe Pro Ser Pro Glu Met Ala Ala Arg Ala His Asp Val Ala Ala Leu
 35 40 45

Ser Ile Lys Gly Ala Ser Ala Ile Leu Asn Phe Pro Asp Leu Ala Gly
 50 55 60

Ser Phe Pro Arg
 65

<210> 27

<211> 68

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> DOMAIN

<222> (1)..(68)

<223> RAP2.1 EREBP-like subclass AP2 domain

<400> 27

Arg Lys Pro Tyr Arg Gly Ile Arg Arg Arg Lys Trp Gly Lys Trp Val
 1 5 10 15

Ala Glu Ile Arg Glu Pro Asn Lys Arg Ser Arg Leu Trp Leu Gly Ser
 20 25 30

Tyr Thr Thr Asp Ile Ala Ala Ala Arg Ala Tyr Asp Val Ala Val Phe
35 40 45

Tyr Leu Arg Gly Pro Ser Ala Arg Leu Asn Phe Pro Asp Leu Leu Leu
50 55 60

Gln Glu Glu Asp
65

<210> 28
<211> 68
<212> PRT
<213> Arabidopsis thaliana

<220>
<221> DOMAIN
<222> (1)..(68)
<223> RAP2.4 EREBP-like subclass AP2 domain

<400> 28
Thr Lys Leu Tyr Arg Gly Val Arg Gln Arg His Trp Gly Lys Trp Val
1 5 10 15

Ala Glu Ile Arg Leu Pro Arg Asn Arg Thr Arg Leu Trp Leu Gly Thr
20 25 30

Phe Asp Thr Ala Glu Glu Ala Ala Leu Ala Tyr Asp Lys Ala Ala Tyr
35 40 45

Lys Leu Arg Gly Asp Phe Ala Arg Leu Asn Phe Pro Asn Leu Arg His
50 55 60

Asn Gly Phe His
65

<210> 29
<211> 66
<212> PRT
<213> Arabidopsis thaliana

<220>
<221> DOMAIN
<222> (1)..(66)
<223> RAP2.8 EREBP-like subclass AP2 domain

<400> 29
Ser Ser Lys Tyr Lys Gly Val Val Pro Gln Pro Asn Gly Arg Trp Gly
1 5 10 15

Ala Gln Ile Tyr Glu Lys His Gln Arg Val Trp Leu Gly Thr Phe Asn
20 25 30

Glu Gln Glu Glu Ala Ala Arg Ser Tyr Asp Ile Ala Ala Cys Arg Phe
35 40 45

Arg Gly Arg Asp Ala Val Val Asn Phe Lys Asn Val Leu Glu Asp Gly
50 55 60

Asp Leu
65

<210> 30
<211> 68
<212> PRT
<213> Arabidopsis thaliana

<220>
<221> DOMAIN
<222> (1)..(68)
<223> RAP2.10 EREBP-like subclass AP2 domain

<400> 30
Asp Lys Pro Tyr Lys Gly Ile Arg Met Arg Lys Trp Gly Lys Trp Val
1 5 10 15

Ala Glu Ile Arg Glu Pro Asn Lys Arg Ser Arg Ile Trp Leu Gly Ser
20 25 30

Tyr Ser Thr Pro Glu Ala Ala Ala Arg Ala Tyr Asp Thr Ala Val Phe
35 40 45

Tyr Leu Arg Gly Pro Ser Ala Arg Leu Asn Phe Pro Glu Leu Leu Ala
50 55 60

Gly Val Thr Val
65

<210> 31
<211> 68
<212> PRT
<213> Arabidopsis thaliana

<220>
<221> DOMAIN
<222> (1)..(68)
<223> RAP2.11 EREBP-like subclass AP2 domain

<400> 31
Lys Thr Lys Phe Val Gly Val Arg Gln Arg Pro Ser Gly Lys Trp Val
1 5 10 15

Ala Glu Ile Lys Asp Thr Thr Gln Lys Ile Arg Met Trp Leu Gly Thr
20 25 30

Phe Glu Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Glu Ala Ala Cys
35 40 45

Leu Leu Arg Gly Ser Asn Thr Arg Thr Asn Phe Ala Asn His Phe Pro
50 55 60

Asn Asn Ser Gln
65

<210> 32
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:EREBP-like
subclass AP2 domain conserved YRG element YRG
motif consensus sequence

<220>
<221> MOD_RES
<222> (4)
<223> Xaa = Val or Ile

<400> 32
Tyr Arg Gly Xaa Arg
1 5

<210> 33
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:EREBP-like
subclass AP2 domain conserved YRG element WAAEIRD
box motif consensus sequence

<220>
<221> MOD_RES
<222> (3)
<223> Xaa = positively charged amino acid

<220>
<221> MOD_RES
<222> (4)
<223> Xaa = Trp, Phe or Tyr

<220>
<221> MOD_RES
<222> (5)
<223> Xaa = Ala or Val

<220>
<221> MOD_RES
<222> (9)
<223> Xaa = Arg or Lys

<220>
<221> MOD_RES
<222> (10)
<223> Xaa = Asp or Glu

<400> 33
Trp Gly Xaa Xaa Xaa Ala Glu Ile Xaa Xaa
1 5 10

<210> 34
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:EREBP-like
 subclass AP2 domain conserved RAYD element
 consensus sequence

<220>
 <221> MOD_RES
 <222> (4)
 <223> Xaa = Thr or Ser

<220>
 <221> MOD_RES
 <222> (5)
 <223> Xaa = Phe or Tyr

<400> 34
 Trp Leu Gly Xaa Xaa
 1 5

<210> 35
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:EREBP-like
 subclass AP2 domain conserved RAYD element RAYD
 consensus sequence

<220>
 <221> MOD_RES
 <222> (5)
 <223> Xaa = positively charged amino acid, Ile or Leu

<400> 35
 Glu Glu Ala Ala Xaa Ala Tyr Asp
 1 5

<210> 36
 <211> 17
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> HELIX
 <222> (1)..(17)
 <223> putative RAP2.7-R1 amphipathic alpha-helix

<400> 36
 Asp Thr Ala His Ala Ala Ala Arg Ala Tyr Asp Arg Ala Ala Ile Lys
 1 5 10 15

Phe

<210> 37
 <211> 16
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> HELIX
 <222> (1)..(16)
 <223> putative ANT-R1 amphipathic alpha-helix

<400> 37
 Met Glu Glu Lys Ala Ala Arg Ala Tyr Asp Leu Ala Ala Leu Lys Tyr
 1 5 10 15

<210> 38
 <211> 18
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> HELIX
 <222> (1)..(18)
 <223> putative RAP2.2 amphipathic alpha-helix

<400> 38
 Asp Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Ala Ala Ala Arg Arg
 1 5 10 15

Ile Arg

<210> 39
 <211> 16
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> HELIX
 <222> (1)..(16)
 <223> putative RAP2.5 amphipathic alpha-helix

<400> 39
 Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Thr Ala Ala Arg Asp Phe
 1 5 10 15

<210> 40
 <211> 18
 <212> PRT
 <213> Arabidopsis thaliana

<220>
 <221> HELIX
 <222> (1)..(18)
 <223> putative RAP2.12 amphipathic alpha-helix

<400> 40
 Lys Thr Ala Glu Glu Ala Ala Arg Ala Tyr Asp Ala Ala Ala Arg Arg
 1 5 10 15

Ile Arg

<210> 41

<211> 16

<212> PRT

<213> Nicotiana tabacum

<220>

<221> HELIX

<222> (1)..(16)

<223> putative EREBP-3 amphipathic alpha-helix

<400> 41

Thr	Ala	Glu	Glu	Ala	Ala	Lys	Ala	Tyr	Asp	Thr	Ala	Ala	Arg	Glu	Phe
1				5					10					15	

<210> 42

<211> 25

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> PEPTIDE

<222> (1)..(25)

<223> AP2 linker region

<400> 42

Lys	Gln	Met	Thr	Asn	Leu	Thr	Lys	Glu	Glu	Phe	Val	His	Val	Leu	Arg
1				5				10						15	

Arg	Gln	Ser	Thr	Gly	Phe	Pro	Arg	Gly
				20				25

<210> 43

<211> 26

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> PEPTIDE

<222> (1)..(26)

<223> ANT linker region

<400> 43

Glu	Asp	Met	Met	Lys	Asn	Met	Thr	Arg	Gln	Glu	Tyr	Val	Ala	His	Leu
1				5				10						15	

Arg	Arg	Lys	Ser	Ser	Gly	Phe	Ser	Arg	Gly
					20				25

<210> 44

<211> 26

<212> PRT

<213> Arabidopsis thaliana

<220>

<221> PEPTIDE

<222> (1)..(26)

<223> RAP2.7 linker region

<400> 44

Met	Lys	Gln	Val	Gln	Asn	Leu	Ser	Lys	Glu	Glu	Phe	Val	His	Ile	Leu
1				5					10					15	

Arg	Arg	Gln	Ser	Thr	Gly	Phe	Ser	Arg	Gly
			20					25	

<210> 45

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:consensus
linker region motif

<220>

<221> MOD_RES

<222> (4)

<223> Xaa = positively charged amino acid

<400> 45

Asn	Leu	Thr	Xaa	Glu	Glu	Phe	Val	His
1			5					

<210> 46

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:consensus
linker region motif

<400> 46

Leu	Arg	Arg	Gln	Ser	Thr	Gly	Phe	Ser	Arg	Gly
1				5					10	

<210> 47

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JOAP2U primer

<400> 47

gttgccgctg ccgtagtg

18

<210> 48

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JOAP2L primer

<400> 48

ggttcatcct gagccgcata tc

22

<210> 49

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.1U
primer

<400> 49

ctcaagaaga agtgcctaac cacg

24

<210> 50

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.1L
primer

<400> 50

gcagaagcta gaagagcgtc ga

22

<210> 51

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.2U
primer

<400> 51

ggaaaatggg ctgcggag

18

<210> 52

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.2L
primer

<400> 52

gttacctcca gcacgaacg ag

22

<210> 53

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.4U
primer

<400> 53

gctggatctt gtttcgctta cg

22

<210> 54

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.4L
primer

<400> 54

gcttcaagct tagcgtcgac tg

22

<210> 55

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.5U
primer

<400> 55

agatgggctt gaaacccgac

20

<210> 56

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.5L
primer

<400> 56

ctggctaggg ctacgcgc

18

<210> 57

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.6U
primer

<400> 57

ttctttgcct cctcaaccat tg

22

<210> 58

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.6L
primer

<400> 58

tctgagttcc aacattttcg gg

22

<210> 59

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.7U
primer

<400> 59

gaaattggta actccggttc cg

22

<210> 60

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.7L
primer

<400> 60

ccattttgct ttggcgcatt ac

22

<210> 61

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.8U
primer

<400> 61

ggcgttacgc ctctaccgg

19

<210> 62

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.8L
primer

<400> 62

cgccgtcttc cagaacgttc

20

<210> 63

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.9U
primer

<400> 63

atcacggatc tggcttggtt c

21

<210> 64

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.9L
primer

<400> 64

gccttcttcc gtatcaacgt cg

22

<210> 65

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.10U
primer

<400> 65

gtcaactccg gcggttacg

19

<210> 66

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.10L
primer

<400> 66

tctccttata tacgccgccg a

21

<210> 67

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.11U
primer

<400> 67

gagaagagca aaggcaacaa gac

23

<210> 68

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.11L
primer

<400> 68

agttgttagg aaaatgggtt gcg

23

<210> 69

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.12U
primer

<400> 69

aaaccattcg ttttcacttc gactc

25

<210> 70

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:JORAP2.12L
primer

<400> 70

tcacagagcg tttctgagaa ttagc

25

<210> 71

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:AP2U primer

<400> 71

atgtgggatc taaacgacgc ac

22

<210> 72

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:AP2L primer

<400> 72

gatcttggtc cacgccgac

19

<210> 73

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.1U primer

<400> 73
aagaggacca tctctcag 18

<210> 74
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:RAP2.1L primer

<400> 74
aacactcgct agcttctc 18

<210> 75
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:RAP2.2U primer

<400> 75
tggttcagca gccaacac 18

<210> 76
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:RAP2.2L primer

<400> 76
caatgcatag agcttgagg 19

<210> 77
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:RAP2.4U primer

<400> 77
acggatttca catcggag 18

<210> 78
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:RAP2.4L primer

<400> 78
ctaagctaga atcgaatcc 19

<210> 79
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.5U primer

<400> 79
taccggtttc gcgcgtag

18

<210> 80
<211> 21
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.5L primer

<400> 80
caccttcgaa atcaacgacc g

21

<210> 81
<211> 22
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.6U primer

<400> 81
ttccccgaaa atgttggaac tc

22

<210> 82
<211> 25
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.6L primer

<400> 82
tgaggagagaa aaaattggta gatcg

25

<210> 83
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.7U primer

<400> 83
cgatggagac gaagactc

18

<210> 84
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.7L primer

<400> 84

gtcggaaaccg gagttacc

18

<210> 85

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.8U primer

<400> 85

tcactcaaag gccgagatc

19

<210> 86

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.8L primer

<400> 86

taacaacatc accggctcg

19

<210> 87

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.9U primer

<400> 87

gtgaaggctt aggaggag

18

<210> 88

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.9L primer

<400> 88

tgcctcatat ggtcagag

19

<210> 89

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.10U primer

<400> 89

tcccggagct tttagccg

18

<210> 90
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.10L primer

<400> 90
caaccggttc caacgatcc

19

<210> 91
<211> 23
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.11U primer

<400> 91
ttcttcacca gaagcagagc atg

23

<210> 92
<211> 24
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.11L primer

<400> 92
ctccattcat tgcatatagg gacg

24

<210> 93
<211> 24
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.12U primer

<400> 93
gctttggttc agaactcgaa catc

24

<210> 94
<211> 22
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.12L primer

<400> 94
aggttgataa acgaacgatg cg

22

<210> 95
<211> 21
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer RISZU 1

<400> 95

ggaytgtggg aaacaagttt a

21

<210> 96

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer RISZU 2

<400> 96

tgcaaagtra cacctctata ctt

23

<210> 97

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer RISZU 3

<400> 97

gcatgwgagc tgtcaaattcc a

21

<210> 98

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer RISZU 4

<400> 98

gaggaagttc vaagtataga

20

<210> 99

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:putative
nuclear localization sequence

<400> 99

Lys Lys Ser Arg

1

<210> 100

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RAP2.3 primer

<400> 100
tcatcgccac gatcaacc

18

<210> 101
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:RAP2.3 primer

<400> 101
agcagtccaa tgcgacgg

18

<210> 102
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:EREBP-like
subclass AP2 domain conserved YRG element WAAEIRD
box motif

<400> 102
Trp Ala Ala Glu Ile Arg Asp
1 5

<210> 103
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:MADS box domain

<400> 103
Met Ala Asp Ser
1

<210> 104
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:AP2-like
subclass AP2 domain conserved YRG element
WEAR/WESH motif

<220>
<221> MOD_RES
<222> (3)
<223> Xaa = Ala or Ser

<220>
<221> MOD_RES
<222> (4)
<223> Xaa = Arg or His

<400> 104

Trp Glu Xaa Xaa

1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/03429

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01H 1/00, 5/00, 5/10; C12N 15/29, 5/04, 5/10

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 800/260, 263, 264, 278, 281, 284, 298, 306; 435/468; 536/23.6; 47/58.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOFUKU et al. Control of Arabidopsis Flower and Seed Development by the Homeotic Gene APETALA2. The Plant Cell. September 1994, Vol. 6, pages 1211-1225, especially page 1215, column 1, lines 3-7. entire document.	44, 45, 47, 51, 52, 58
A	OKAMURO et al. The AP2 domain of APETALA2 defines a large new family of DNA binding proteins in Arabidopsis. Proceedings of the National Academy of Sciences USA. June 1997, Vol. 94, pages 7076-7081.	1-59

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*N* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*X* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

27 MAY 1999

Date of mailing of the international search report

14 JUN 1999
 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer


 MARY MOSHER

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/03429

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

800/260, 263, 264, 278, 281, 284, 298, 306; 435/468; 536/23.6; 47/58.1

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, Dialog files 351, 5, 50. Search terms: APETALA?, AP2, AP 2, PLANT#, SEED#, MASS, SIZE, WEIGHT, LARGE, LARGER, INCREAS?, SMALL, SMALLER, RECOMBIN?, TRANSGENIC?, AU=JOFUKU ?, ROL, ROLC, CLON?, SEQUENC?, AINTEGUMENT?